

# Temperature-Mediated Alterations of the Plant Apoplast as a Mechanism of Intracellular Freezing Stress Avoidance

A Thesis Submitted to the College of Graduate Studies and Research  
in Partial Fulfillment of the Requirements  
for the Degree of Masters of Science  
in the Department of Plant Sciences  
University of Saskatchewan  
Saskatoon

By

**Jun Liu**

© Copyright Jun Liu November 2015. All rights reserved.

## PERMISSION TO USE

In presenting this thesis in partial fulfilment of the requirements for a Postgraduate degree from the University of Saskatchewan, I agree that the Libraries of this University may make it freely available for inspection. I further agree that permission for copying of this thesis in any manner, in whole or in part, for scholarly purposes may be granted by the professor or professors who supervised my thesis work or, in their absence, by the Head of the Department or the Dean of the College in which my thesis work was done. It is understood that any copying or publication or use of this thesis or parts thereof for financial gain shall not be allowed without my written permission. It is also understood that due recognition shall be given to me and to the University of Saskatchewan in any scholarly use which may be made of any material in my thesis.

Requests for permission to copy or to make other use of material in this thesis in whole or part should be addressed to:

Head of the Department of Plant Sciences  
College of Agriculture and Bioresources  
51 Campus Drive  
University of Saskatchewan  
Saskatoon, Saskatchewan S7N 5A8, Canada

## ABSTRACT

Cold hardy Japanese bunching onion (*Allium fistulosum* L.) was used as a novel model system to examine the role of the apoplast in intracellular freezing avoidance. Since intracellular freezing avoidance is critical to both sensitive and resistant plants, non-acclimated and cold acclimated onion tissue was compared. The large 250 (length) x 50 (width) x 90  $\mu\text{m}$  (thickness) intact single cell epidermal layer system allowed direct observation of the freezing process during freezing and thawing on a single cell basis in live intact tissues in non-acclimated (23/18°C) and acclimated (2 weeks at 12/4°C) plants. Under acclimation treatment, freezing resistance increased from an  $\text{LT}_{50}$  of -12°C at zero time to -27°C after two weeks exposure. Ice nucleation was always initiated in the apoplastic space in both non-acclimated and acclimated cells and then penetrated into the intracellular space. We provide direct evidence that lethal intracellular freezing damage was avoided in cold acclimated and  $\text{CaCl}_2$  treated cells through visible blockage of ice propagation from the apoplastic space to the intracellular region. Subsequent investigation of the apoplastic space revealed cold acclimation increased: a) the insoluble crude cell wall and pectin (galacturonic acid) content, b) total cell wall protein quantity, c) pectin methylesterase activity in the apoplastic space particularly in epidermal cells compared to non-epidermal cells, and d) the levels of un-methylated pectin. The increasing freezing resistance of Japanese bunching onion more depends on the efficiency of PME not only the increase in the amount of apoplast compositions. Collectively, these changes likely led to the observed significant reduction in cell wall permeability. In this way, freezing stress resistance may potentially be increased through avoidance of intracellular freezing by altering key factors related to apoplast permeability and blocking ice propagation across the cell wall.

## ACKNOWLEDGEMENTS

I would like to express my sincerest appreciation to my supervisor Dr. K.K. Tanino for her kindly assistance, support and motivation through my graduate study and research. Her vast knowledge, serious attitude and commitment to research made this study a success. I want to extend my sincere thanks to all my committee members, Dr. Y. Bai, Dr. G. Gray, Dr. M. Båga and Dr. Y. Wei. Their questions, suggestions, encouragements and comments guided me to seek answers in new perspectives. My special thanks go out to Dr. H. Li and Dr. P. Vijayan for their great endeavors for instructing my experiments. I am also grateful to my external examiner.

My thanks also go to E. Siemens at the Agriculture Greenhouse for helping me throughout my greenhouse experiment. My special thanks are extended to Dr. Y. Lee who taught me to do sectioning and immunolabeling and Trine Hvoslef-Eide who introduced the microscopy resources in Norwegian University of Life Sciences, Dr. G. Liu in the Department of Biology for providing and guiding me using the equipment for my experiment, Ricky Lee and Dr. Supratim Ghosh in Department of Food and Bioproduct Sciences for helping me to figure out the problems in the measurements of cryostage behavior. My sincere appreciation is extended to G. Swerhone and J. Lawrence in National Hydrology Research Institute, F. Borondics and X. Liu in Canadian Light Source for measuring my sample and providing me the valuable results. Diligent assistance from my lab mates, namely J. Xu and P. Banik are highly appreciated as well.

Financial support provided through China Scholarship Council and University of Saskatchewan has made my research and life much easier over the last few years. I am grateful to have received the Vandeveld, Rene Bursary in Plant Sciences. I am also grateful for the travel awards received through the University Postgraduate Scholarships for my learning experience in Europe.

Last but not least, I would like to thank my family members and friends, especially my husband Lei Ren, who supported and encouraged me through it all.

# TABLE OF CONTENTS

	<u>page</u>
PERMISSION TO USE .....	i
ABSTRACT.....	ii
ACKNOWLEDGEMENTS .....	iii
TABLE OF CONTENTS.....	iv
LIST OF TABLES .....	vi
LIST OF FIGURES .....	vii
1.0 INTRODUCTION .....	1
2.0 LITERATURE REVIEW .....	5
2.1 Freezing Stress .....	5
2.1.1 Freezing Process.....	5
2.1.2 Freezing Avoidance and Freezing Tolerance .....	9
2.2 Apoplast.....	12
2.2.1 The Structure of the Apoplast.....	12
2.2.2 Function of the Apoplast .....	16
2.2.3 Responses to Cold Acclimation and Role in Freezing Resistance .....	17
2.3 Calcium.....	21
2.3.1 Calcium is an Essential Element in Plants.....	21
2.3.2 Role of Calcium in the Apoplast and Freezing Stress Resistance .....	22
2.4 <i>Allium fistulosum</i> L. ....	23
3.0 MATERIALS AND METHODS .....	25
3.1 Plant Materials and Growth .....	25
3.2 Freezing Stress Treatments .....	25
3.3 Analysis of Freezing Process .....	27
3.4 Study of Apoplast Permeability .....	27
3.5 Analysis of Pectin Methyl Esterification .....	28
3.5.1 Sample Preparation and Fixation.....	28
3.5.2 Analysis of Pectin Methyl Esterification by Immunohistochemistry .....	29
3.5.3 Determination of PME Activity .....	29
3.6 Cell Wall Compositional Analysis.....	30
3.6.1 Cell wall isolation.....	30
3.6.2 Matrix Polysaccharide composition .....	31
4.0 RESULTS AND DISCUSSION .....	33
4.1 Freezing Stress Resistance.....	33
4.2 Analysis of Freezing Process .....	36
4.3 Studies of Apoplastic Permeability.....	41
4.4 Analysis of The Degree of Pectin Methyl-esterification.....	44
4.5 Cell Wall Compositional Analysis.....	46
4.5.1 Determination of Cell Wall Quantity in Ligule Tissue .....	46

4.5.2	Cell Wall Carbohydrate Analysis of Ligules.....	47
4.6	PME Activity in Epidermal Tissues of Ligules .....	48
4.6.1	Determination of Total Cell Wall Protein Content.....	48
4.6.2	PME Activity Analyses .....	50
5.0	CONCLUSIONS .....	58
6.0	FUTURE WORK AND SIGNIFICANCE .....	60
7.0	REFERENCES .....	62
APPENDIX A: Growth and morphology of <i>Allium fistulosum</i> L.....		93
APPENDIX B: The cryostage microscope equipment .....		94
APPENDIX C: Protoplastic streaming and ice penetration in the cells.....		95
APPENDIX D: Analysis of freezing behavior in treated epidermal layer cells of ligules .....		96
APPENDIX E: Confocal microscope and fluorescein penetration.....		98
APPENDIX F: Evos digital inverted microscope (Department of Plant Sciences) .....		99
APPENDIX G: Analysis of pectin methyl esterification in ligule epidermal cells .....		100

## LIST OF TABLES

<u>Table</u>	<u>Page</u>
Table 4.1 Summary of extracellular and intracellular freezing temperature in NA, ACC and NA+Ca treated plants.....	37

## LIST OF FIGURES

<u>Figure</u>	<u>page</u>
Figure 2.1 The freezing effect on plant cells.....	6
Figure 2.2 Schematic structural models of type I and type II walls.....	13
Figure 2.3 The formation of pectin networks by covalent and ionic bonds.....	14
Figure 3.1 The experimental area of Japanese bunching onion.....	26
Figure 3.2 The ligule and collar region of the plant.....	27
Figure 4.1 Effect of cold acclimation time on freezing resistance.....	34
Figure 4.2 Effect of cold and calcium on freezing hardness.....	35
Figure 4.3 Organization of the plant cuticle.....	39
Figure 4.4 The impact of acclimation on apoplast permeability.....	42
Figure 4.5 The diffusion rate of fluorescein in NA and ACC.....	43
Figure 4.6 Immunohistochemical studies of pectin composition in NA and ACC.....	45
Figure 4.7 The relative content of onion crude cell wall mass during cold acclimation.....	47
Figure 4.8 Cell wall components of the epidermal layer.....	48
Figure 4.9 Relative total cell wall protein content during cold acclimation.....	49
Figure 4.10 Relative PME specific activity during cold acclimation.....	51
Figure 4.11 The PME activity in the whole sheath of onion species.....	53



## 1.0 INTRODUCTION

Temperature has had a primary influence on adaptation throughout plant evolutionary history. Future temperatures are predicted to increase under global climate change with night temperatures increasing more than day temperatures (Easterling and Horton, 1997; IPCC, 2007). According to Gu *et al.* (2008) and as reviewed in Storey and Tanino (2012), injury due to low temperature stress will become even more important under global warming, not less. Since temperature also influences the ability to increase freezing resistance through the process of cold acclimation, the ability to acclimate in response to temperature is an important survival mechanism in temperate plants.

Kozlowski and Pallardy (1997) divided cold acclimation into two sequential stages: Stage I: The accumulation of carbohydrates and lipids in the autumn, which are substrates and energy sources for the next metabolic phases; Stage II: Freezing temperatures to synthesize proteins and membrane lipids and change structures to acclimate against cold winters. Cold acclimation is regulated by both low temperature and photoperiod and shares similarities with the important environmental induction parameters of growth cessation and dormancy (Weiser, 1970).

Water diffusivity (Zeuthen, 2001; Welling and Palva, 2006; Yooyongwech, 2008; Kalcsits *et al.*, 2009) and carbohydrate accumulation in the cell wall (Olien and Clark, 1993; Livingston *et al.*, 2009; Tanino *et al.*, 1990, 1991, 2010) are significantly connected with the development of low temperature stress resistance. The regulation of water flow across the cell wall is largely attributed to aquaporins (Carvajal *et al.*, 1998) and plasmodesmata (Kuo *et al.*, 1974; Anisimov and Egorov, 2002; Liesche *et al.*, 2011). Aquaporin genes are downregulated in the transition periods during dormancy phases (Mazzitelli *et al.*, 2007). Most of the environmental stress factors, including low temperature, can stimulate the generation of intracellular signals and depositions of callose at plasmodesmata during dormancy (Sivaguru *et al.*, 2000; Rinne *et al.*, 2011; van der Schoot *et al.*, 2011), thus reducing water flow into the cells. However, the role of the apoplastic matrix itself (cell wall and middle lamella) during cold stress has not received much attention.

The apoplast is a physical barrier against environmental intrusions, including pathogens and abiotic factors, *i.e.* ice propagation and dehydration stress through excessive water loss.

Plant species in the temperate zones are vulnerable to natural low temperatures and cold hardy plants first freeze outside the plasma membrane in the extracellular space of the apoplast (cell wall, middle lamella) (Levitt, 1980). Water inside the cells will then migrate out to freeze in the apoplast, following its vapour pressure deficit, thus leading to cellular dehydration and osmotic shrinkage of the cell (Dowgert and Steponkus, 1984; Ruelland *et al.*, 2009). To a certain degree, the mechanical characteristics of cell walls appear to adjust to the kinetics and the range of dehydration being stimulated by freezing, thereby improving the ability of plants to resist extracellular freezing (Solecka *et al.*, 2008). The freezing process can result in cell dehydration stress. As the temperature is lowered, the more serious cell dehydration becomes (Pearce and Ashworth, 1992). However, dehydration resulting from extracellular freezing can help cells to avoid intracellular freezing and increase survival at very low, even liquid-nitrogen temperatures (Sakai and Larcher, 1987). Conversely, if intracellular freezing occurs, the cells will die (Levitt, 1980). Thus, the apoplast matrix represents a common barrier to both water loss out from the cell (freezing tolerance), as well as a barrier to ice penetrating into the intracellular space (freezing avoidance).

Gusta *et al.* (2009) indicated there are seven different freezing types in plants, such as partially or fully cold acclimation, and supercooling or deep supercooling. However, regardless of the freezing process, the common critical factor to frost survival is avoidance of intracellular freezing. Since the apoplast represents the first barrier to ice, the changes within the apoplast under cold acclimation will be examined in this thesis. While several studies have examined modification of ice crystals within the apoplastic space by carbohydrates (Griffith *et al.*, 1985; Olien and Lester, 1985; Olien and Clark, 1993), this thesis focuses on the apoplast as a barrier preventing ice propagation across the cell wall to freeze the intracellular space.

The principal elements in the apoplast (consisting of middle lamella and cell wall) are polysaccharides consisting of monosaccharides. The cell wall fraction is a biphasic structure consisting of cellulose microfibrils held together by a gel-like matrix which constitutes about two-thirds of the cell wall's dry weight and is composed of cross-linking extensin glycoproteins and non-cellulosic polysaccharides including pectins and hemicelluloses (Fry, 1986; Pelloux *et al.*, 2007). Other apoplastic components are lignin, suberin, wax, cutin and water. The middle lamella is primarily composed of pectins. Pectins are chiefly divided into xylogalacturonan, homogalacturons (HGA), rhamnogalacturonan I (RGI) and rhamnogalacturonan II (RGII). These

components serve many functions including determining apoplast porosity, providing a charged wall surface supporting cell-cell adhesion (i.e., middle lamella), cell-cell identification, pathogen recognition and other roles.

The gelation and rigidity of the cell wall is regulated by pectin substances through pectin methylesterase (PME) (Willats *et al.*, 2001). PME catalyzes the de-esterification of pectin methyl groups, permitting  $\text{Ca}^{2+}$  binding to the carboxylate ions and generating calcium cross-linkages (Pelloux *et al.*, 2007). In poplar cambium, acidic pectins and calcium causes early stiffening of maturing phloem cells (Guglielmino *et al.*, 1997a) and apparent changes in PME isoforms (Guglielmino *et al.*, 1997b). Cell wall-bound PME's are temperature- sensitive (Anthon and Barrett, 2006) but most of this evidence comes from the food processing literature and few such studies have been done on living plants. Wisniewski and Davis (1995) illustrated deep supercooling in trees is associated with pectins preventing water loss and ice nucleation within the cells. A subsequent study (Solecka *et al.*, 2008) demonstrated cold acclimation of winter oil-seed rape is increased by the temperature-dependent PME. Although it is well-known that the mediation of water movement is critical to plant growth and development, there are few studies examining the role of the cell wall, and pectins in particular, in regulating water flow and ice propagation (Wisniewski and Davis, 2005). Most studies of the apoplast water migration have been focused on aquaporins (Mazzitelli *et al.*, 2007) and plasmodesmata (Liesche *et al.*, 2011).

Previous studies have shown that Japanese bunching onion is a suitable system for studying changes in the structure and components of the cell wall in response to intracellular freezing avoidance (Tanino *et al.*, 2013). Japanese bunching onion is a very hardy species and can tolerate extreme environmental stress. The advantage of this plant for apoplast studies is an easy to peel intact single epidermal layer devoid of chlorophyll that facilitates immunofluorescence studies. In addition, the cells are large (250 x 50 x 90 $\mu\text{m}$ ). The plant acclimates quickly to cold (increases in freezing resistance from -12°C to -27°C in two weeks of acclimation). All of these advantages make Japanese bunching onion an ideal system to study freezing stress mechanisms.

## 1.1 Hypothesis

Reduced permeability of the apoplast through pectin modification will increase freezing avoidance and freezing stress survival.

## **1.2 Objectives**

- 1) To determine if the permeability of the apoplast (cell wall/middle lamella) plays a role in freezing avoidance.
- 2) To determine if cold acclimation decreases apoplastic permeability through apoplast compositional changes.

## **2.0 LITERATURE REVIEW**

### **2.1 Freezing Stress**

Plant stress can be divided into two types: biotic stress and abiotic stress. Biotic stress involves plant damage from other living organisms, such as bacteria, fungi, and insects. Abiotic stress, also known as environmental stress, is referred to as the adverse influences on the existing organisms resulting from non-living factors (Ortbauer, 2013). The non-living factors include: extreme temperatures, drought, flood, salt, wind, etc. Each year, millions of tonnes of potential crops and other plants yield fail to materialize because of various abiotic stress factors that impact the growth and maturation of the plants (Xiong and Zhu, 2001; Kozłowski, 2002). Plants are commonly very susceptible to environmental stimuli. At the same time, plants can physiologically and structurally acclimate and adjust in order to survive in their environment.

Based on the research of Intergovernmental Panel on Climate Change (IPCC, 2014), the extreme temperatures have been changing for decades. It is predicted that the global average surface temperature may rise from 2.5 to 7.8°C in 2100 (IPCC, 2014). Increasing temperature has a negative effect on agriculture because of the frequency of drought, pests and diseases which may arise. There will be fewer cold days and cold nights and frosts, more hot days and hot nights and heat waves. In addition, snow and ice covers are decreasing widely. Snow and ice plays a vital role in the global climate system and normally provides the avoidance protection for plants. The growing season will also be extended in which spring occurs earlier and autumn later. This has already increased damage due to early spring bloom which is more susceptible to frost damage. In 2007, over \$2 billion US damage occurred to frost sensitive horticulture crops (e.g. bedding plants, blossoms of pear, apple and peach) in the USA over a two day weekend (Gu *et al.*, 2008). Sensitive plants only have one mechanism of freezing stress resistance and that is frost avoidance. Therefore, it will become increasingly critical to understand the mechanism of frost avoidance in plants.

#### **2.1.1 Freezing Process**

One of the most significant stresses is low temperature which includes two types of cold stress: chilling stress and freezing stress. Chilling stress is commonly defined to occur at low temperatures but above zero (from 0 to 10°C) without ice formation, while freezing stress is

exerted at negative temperatures with ice nucleation in plant tissues (Thomashow, 1999; Kolaksazov *et al.*, 2013). Freezing stress can affect the growth of aerial parts of plants (such as leaf expansion), causing wilting, chlorosis and necrosis (Mahajan and Tuteja, 2005; Sanghera *et al.*, 2011), alter root growth (Noh *et al.*, 2009), and even result in cell collapse and death (Stefanowska *et al.*, 1999; Mahajan and Tuteja, 2005). The freezing effect on plant cells is described in Figure 2.1. Freezing injury is thought to occur primarily in the plant plasma membrane mainly due to the severe dehydration induced by freezing (Pearce and Willison, 1985; Steponkus *et al.*, 1998; Pearce, 2001; Thomashow, 2001). Once the ice crystals penetrate the symplast or grow directly into the intracellular spaces, it is immediately fatal since the intracellular ice can physically destroy the structures of plasma membrane and the integrity of the living protoplasm and this process is irreversible (Burke *et al.*, 1976; Gusta *et al.*, 2004).




Figure 2.1 The freezing effect on plant cells (Ruelland *et al.*, 2009).

Freezing stress in plants can be sorted into at least seven different types (Stushnoff *et al.*, 1984; Gusta *et al.*, 2009). The first freezing type includes the succulent tender tissues like tomatoes and corn containing high moisture content such as tropical plants. In this pattern, the plants are very sensitive to both chilling and frost and the injury occurs just below freezing from -2°C to -5°C. The second freezing type appears in partially acclimated tissues or mature tissues with its LT<sub>50</sub> values from -3 °C to -10 °C. The third freezing pattern occurs in fully cold-acclimated hardy species with low water content (< 75% water) such as winter cereals and trees that do not supercool, and for these species the freezing injury is mainly due to extracellular freeze-induced dehydration. The fourth freezing type is related to the ability of certain cells (such as xylem ray parenchyma cells) to supercool. When the cooling rate of the tissue is greater than 3 °C/hour, the fifth freezing type occurs, which is not regarded as true supercooling. This type

occurs when the low temperature exotherm (LTE) disappears and the temperature of the LTE does not correspond to the killing temperature in slowly cooled seeds. The sixth freezing type is related to differential organ supercooling and extraorgan freezing in non-acclimated cereals in which contains barriers to prevent ice penetration. The seventh freezing type involves vitrification or glass formation which occurs at sub-zero temperatures when water is converted into a glass-like amorphous solid but does not form the crystalline structure.

At the cellular level, ice crystals in freezing resistant plants are first generated on the surface of cell walls and spread throughout the extracellular space (Burke *et al.*, 1976; Guy, 1990). Ice can be limited to the external regions of the cell by the intact cell wall and plasma membrane and a slow cooling rate. This process is called extracellular freezing (Levitt, 1980; Guy, 1990). The intracellular water diffuses down its vapour pressure gradient caused by extracellular freezing, to form ice crystals in the apoplastic space resulting in cell dehydration and cell osmolarity, induces cell shrinkage and finally cell collapse (Ashworth *et al.*, 1988; Yelenosky and Guy, 1989; Guy, 2003). Accompanying the falling temperatures, more intracellular water diffuses out and the extent of dehydration increases (Johansson and Krull, 1970; Gusta *et al.*, 1975; Ashworth and Pearce, 2002). Once the dehydration exceeds the critical range, the cells are killed by a combination of low temperature and desiccation stress.

The other freezing condition is termed intracellular freezing which may occur due to ice nucleation throughout the protoplast and vacuole (Levitt and Scarth, 1936) or the penetration to the intracellular spaces via the extracellular ice (Mazur, 1969; Steponkus *et al.*, 1983) when the cooling rate is rapid ( $\geq 3^{\circ}\text{C}/\text{hour}$ ). The accumulation of ice in the middle lamella can potentially lead to cell disruption partly due to the ice adhesions between the cells, cell walls and membranes (Levitt, 1980). The direct consequence of intracellular freezing is lethal to all the cells, which may be related to the physical destruction of the plasma membrane due to internal ice crystal propagation (Muldrew *et al.*, 2004). However, even though the cooling rate is slow ( $< 1^{\circ}\text{C}/\text{hour}$ ), there is an exception—intracellular freezing may not take place but deep supercooling occurs in some tissues such as xylem (Wisniewski and Davis, 1995), bark (Quamme, 1985) and flower buds (Sakai, 1978; Ashworth, 1984). Supercooling is the process where a liquid or a gas does not become a solid state below its freezing point when lowering the temperature (Moore and Molinero, 2011).

The freezing process in plants can be driven by: water potential in both extracellular and intracellular water (Wenkert, 1980; Levitt, 1972, 1980), all intact effective cell wall and plasma membrane serving as a freezing barrier to avoid intracellular freezing (Dowgert and Steponkus, 1984; Steponkus *et al.*, 1983; Levitt and Scarth, 1936; Steponkus *et al.*, 1982), the existence of heterogeneous nucleators (Burke *et al.*, 1976; Sakai and Larcher, 1987), and the rate of cooling (Mazur, 1963, 1969; Steponkus *et al.*, 1983; Guy, 1990). Ice formation in plant tissues is triggered in nucleation sites where water molecules come together to form an ice nucleus when exposed to low temperature (Burke and Lindow, 1990). This process belongs to heterogeneous nucleation which can be induced by other substances including ice nucleation-active (INA) bacteria, the biological molecules and structures, organic and inorganic debris (Pearce, 2001). The nucleation sites are usually of a specific size and shape, and are related to the composition of the cell wall. Both composition and structure of the cell wall can assist the plant tissues to control the ice nucleation sites. Ice formation is thought to initially occur in these areas with the least negative osmotic potential such as the large vessels of xylem of leaves and stems (Levitt, 1980; Sakai and Larcher, 1987), thereafter spread rapidly into the extracellular spaces of other tissues. Recent evidence indicates ice nucleation can initiate in the phloem and cortical tissue of *Rhododendron japonicum* and *Cornus officinalis* (Ishikawa and Sakai, 1982), stems of blueberry (Kishioto *et al.*, 2014) and *Rhododendron japonicum* flower bud scales (Ishikawa *et al.*, 2015).

Due to their immobility, plants have to sense and respond to the fluctuating environmental temperature. One of the amazing abilities of plants is to increase their freezing resistance through pre-exposure to low but non-freezing temperatures, which is called cold acclimation (Levitt, 1980; Sakai and Larcher, 1987; Guy, 1990; Steponkus, 1990; Thomashow, 1999). Numerous alterations in morphology, metabolism, physiology, and gene expression occur during cold acclimation (Thomashow, 1999; Levitt, 1980; Xin and Browse, 1998) in order to increase the possibility of plant survival under low temperature and/or freeze-induced dehydration stress. With these alterations, plants can tolerate the extracellular dehydration stress induced by extracellular freezing (freezing tolerance) and/or avoid intracellular ice nucleation (freezing avoidance). The multiple mechanisms of plant resistance usually do not exist independently but occur in the same organs to defense freezing injury. For example, the flower buds of *Rhododendron japonicum* avoid freezing damage by supercooling (freezing avoidance) (Ishikawa and Sakai, 1981) while the flower buds of *Rhododendron japonicum* can tolerate the



accumulation of ice in a particular space outside the organ such as bud scale which can be regarded as a barrier through partial dehydration (freezing tolerance) (Ishikawa and Sakai, 1982). In addition, the existence of barriers such as the cell wall and plasma membrane can help to isolate the cells from extracellular freezing and inhibit the loss of water molecules to the surrounding ice.

### **2.1.2 Freezing Avoidance and Freezing Tolerance**

There are two major categories of freezing resistance: freezing avoidance and freezing tolerance (Levitt, 1972, 1980). Freezing avoidance can occur when plants affect the freezing process by preventing the penetration of ice crystals into the tissue or the cellular levels. Examples include deep supercooling of internal tissue water which helps to avoid cell dehydration caused by extracellular ice formation (Ishikawa and Sakai, 1981; Wisniewski *et al.*, 1987) and extraorgan freezing (Ishikawa and Sakai, 1982; Fujikawa *et al.*, 2000, 2009; Kuroda *et al.*, 2003; Endoh *et al.*, 2009). In addition, freezing avoidance can also include shedding or insulating vulnerable plant parts (Raulston and Tripp, 1994). For example, alpine plants warm their shoots by burying themselves below ground and thereby avoiding severe freezing stress, while permitting leaves to freeze independently in the air (Körner, 2003). In this way, separation of ice nucleation and the prevention of ice propagation from the frozen shoot enable susceptible reproductive structures to avoid freezing stress (Hacker *et al.*, 2011).

Freezing avoidance regulates the nucleating position of ice crystals in the plant (Wisniewski *et al.*, 2014). Heterogeneous ice nucleation, triggered by ice nucleation active substance (INAS) including inorganic substances such as silver iodide (Vonnegut, 1947) and organic substances such as ice-nucleating-active bacteria (e.g. *Pseudomonas syringae*) (Ashworth and Kieft, 1995) is different from homogeneous ice nucleation, in which ice form autonomously (Wisniewski *et al.*, 2014). Silver iodide (AgI) is an effective nucleus agent to form the ice crystals since the crystal structure (the dimensions of the unit) of AgI is similar to ice (Vonnegut, 1947). Therefore, the application of AgI is easy to initiate ice nucleation in cellular space. Occurrence of extrinsic or intrinsic ice nucleation depends on the presence of INAS on the plant surface or within the plant (Pearce, 2001). However, INAS are not the exclusive trigger for extrinsic ice nucleation. Wettability of plants plays an important role too. Reducing leaf wettability leads to the reduction of extrinsic ice nucleation in potatoes, grapevine, lemon and tomato (Wisniewski *et al.*, 2002). Intrinsic ice nucleation is regulated by both INAS and anti-ice

nucleating substances, such as antifreeze proteins (AFPs) and fructans. While AFPs are regarded to play a crucial role in affecting the shape and size of ice crystals (Griffith *et al.*, 2005), fructans are thought to stabilize cell membranes and change the morphology and growth pattern of ice crystals (Olien and Marchetti, 1976; Valluru and Van den Ende, 2008). As one of the most well-known examples of freezing avoidance is supercooling, this phenomenon allows water in plant cells to remain in a liquid phase and avoid freezing at low, subfreezing temperatures (Wisniewski *et al.*, 2014) due to solutes within the cell solution reducing the freezing point. In addition, anti-nucleating substances such as tannins and related polyphenols and glycosides can also enhance supercooling extent (Kuwabara *et al.*, 2013; Kasuga *et al.*, 2010). Due to deep supercooling, apoplastic water freezing of xylem ray parenchyma cells can be avoided (George *et al.*, 1982). The intracellular space does not lose water even in the presence of ice in apoplastic spaces via deep supercooling (Fujikawa *et al.*, 1994), thereby maintaining the original volumes. Extraorgan freezing is freezing of large extracellular cavities such as bud scales which serve as preferred freezing sites in tree buds (Sakai and Larcher, 1987). Ice propagation is prevented in the dry region created by water withdrawal to the extracellular cavities.

Once intracellular freezing occurs, it is lethal for all biological materials (Levitt, 1980). Therefore, different types of barriers exist in plants, isolating the supercooled cells from external nucleation in order to block the ice propagation and prevent water loss to surrounding ice. Thermal ice barriers refer to plant structures that are thermally buffered to separate them from those directly exposed to freezing stress. Structural ice barriers are associated with compositional alteration of the plasma membrane and cell wall architecture to inhibit ice propagation (Ashworth and Abeles, 1984; Steponkus, 1984; Uemura and Steponkus, 1997). For instance, a break in the epicuticular layer provides a pathway for ice to physically propagate into the leaf. Hence, thick cuticles serve as effective structural ice barriers. Supercooled cell walls are generally thicker and less flexible (Fujikawa *et al.*, 1999). The xylem ray parenchyma cells including pit membrane in woody plants (Wisniewski and Davis, 1989; Wisniewski *et al.*, 1991b), and flower buds in many *Prunus* species and other plants (Quamme, 1974; Graham and Mullin, 1976; Biermann *et al.*, 1979; Wolpert and Howell, 1984; Sakai *et al.*, 1986) are also regarded as ice barriers resulting from physical characteristics to prevent ice propagation and water loss (Chalker-Scott, 1992). Through use of Infrared Video Thermography (IRVT) it is known that ice in cranberry stems cannot propagate into the fruit (Workmaster *et al.*, 1999). Ice

barriers are also present in the crowns (Single and Marcellos, 1981) and at nodes (Zámečník *et al.*, 1994; Wisniewski *et al.*, 1997; Pearce and Fuller, 2001) of cereals and beans, respectively, where they delay the spread of ice throughout the tissue.

Freezing tolerance is defined as the allowance of ice nucleation but the strain is decreased or eliminated through resistance to extracellular freezing dehydration and cell shrinkage. Thus, freezing tolerance refers to the capability of plants to tolerate the presence of ice in their extracellular tissues and the concomitant dehydration stress through cold acclimation (Gusta and Wisniewski, 2013). The tolerance generally associated with biochemical adaptations and alterations in gene regulation. Freezing injury in plants is partly due to interruption of plant water status when cellular dehydration appears because of ice generation outside the cell in the freezing process (Ashworth and Kieft, 1995; Verslues *et al.*, 2006). Freezing resistance mechanisms could be referred to as dehydration resistance mechanisms which provide protection to cellular architecture from the impacts of dehydration. There are various mechanisms of dehydration or freezing resistance, including: (1) protection and repair mechanisms which are a primary means to tolerate desiccation stress induced by freezing and involve late-embryogenesis-abundant (LEA) (Close, 1997) and dehydrin proteins (COR15), compatible solutes (Hinch and Hagemann, 2004), ROS scavenging (Apel and Hirt, 2004); (2) a collection of solutes and cell wall hardening, which is associated with dehydration avoidance mechanisms (Verslues *et al.*, 2006); (3) accumulation of hormones such as ABA, which is a key regulatory factor in reducing dehydration stress via regulating stomatal conductance (Schroeder *et al.*, 2001) and accumulating compatible solutes (Ober and Sharp, 1994); (4) the changes of plasma membrane organization and functionality such as lipids (e.g. increase in the proportion of phospholipids) (Uemura *et al.*, 2006) and proteins (e.g. dehydrin family proteins) (Kosová *et al.*, 2007) is a critical prerequisite in order to survive from dehydration stress.

Expression of several different types of proteins, including dehydrins (Hanin *et al.*, 2011; Hara, 2010), Cold Responsive proteins (Gilmour *et al.*, 1992), increase under low temperature condition. Although not conclusive, these proteins are assumed to be associated with freezing tolerance, since they can protect cellular components from denaturation (Burchett *et al.*, 2006) by interacting with sugars to inhibit desiccation (Wolkers *et al.*, 2001).

Plants do not gain freezing tolerance capability effectively without the accumulation of sugars (Hurry *et al.*, 1994; Dahal *et al.*, 2012). Huge differences in cold acclimation capability

between spring (e.g. LT50, -9 °C) and winter cereals (e.g. LT50, -25 °C) appears to be due to the inability of spring cereals to effectively accumulate sugars and anthocyanins at low non-freezing temperatures, unlike winter cereals (Hu *et al.*, 2001; Hüner *et al.*, 1993; Dahal *et al.*, 2012).


## **2.2 Apoplast**

### **2.2.1 The Structure of the Apoplast**

The apoplast is a free diffusional compartment outside the plasma membrane and includes the walls of living cells and entire nonliving cells, middle lamella, intercellular air spaces, water and solutes (Peterson and Cholewa, 1998; Rose, 2003). The cell wall consists of a complex matrix of polysaccharides and other polymers, and plays a crucial role in plant cell survival. Although appearance and composition of apoplast vary in different cell types (Albersheim *et al.*, 2011), basically they are composed of three layers: middle lamella, primary walls and secondary walls (Buchanan *et al.*, 2000). Primary cell walls appear in the young cells of growing plant tissues, and mainly consist of cellulose, hemicellulose and pectin (McNeil *et al.*, 1984). Secondary cell walls form after the cessation of cell growth; they are highly different in structure and composition compared with primary cell walls, containing cellulose, xylan and lignin. Generally, secondary cell walls are thicker and have more strength than primary cell walls due to the presence of lignin (Vogel, 2008). The middle lamella is a thin layer of material connecting the walls of adjacent cells and is composed almost entirely of pectin. Plant cell walls also contain four types of structural proteins: hydroxyproline-rich glycoproteins (HRGP), arabinogalactan proteins (AGP), glycine-rich proteins (GRPs), and proline-rich proteins (PRPs) (Rose, 2003). In addition, cell walls contain abundant enzymes such as hydrolases and esterases. Cell walls from the epidermis and endodermis may contain higher levels of suberin or cutin than other cells.


The primary cell walls of angiosperms are classified into two distinct types as illustrated in Figure 2.2 (Carpita and Gibeaut, 1993; Yokoyama and Nishitani, 2004; Jones *et al.*, 2005; Vogel, 2008). Type I cell walls exist in all dicots and non-graminaceous monocots such as aroids and lilioids and is a network of cellulose microfibrils interlocked with xyloglucan (XyG) and structural proteins and embedded in a matrix of pectic polysaccharides. Type II cell walls are found in the Poaceae and some related monocot families such as rushes and gingers. They consist of similar cellulose microfibril structures as the Type I cell walls but with a framework

interlocked with glucuronoarabinoxylans (GAX). The amount of pectin and structural proteins in the cell wall can be used to distinguish these two types of primary cell walls. Type I cell walls have 20-35% pectin (O'Neil and York, 2003) and 10% structural proteins (O'Neil and York, 2003; Zablackis *et al.*, 1995), while Type II cell walls have 5% pectin (Ishii, 1997) and 1% structural proteins (O'Neil and York, 2003). Based on these characteristics, the cell walls of onion (*Allium cepa*) belong to the Type I cell wall (McCann *et al.*, 1992; Carpita and Gibeaut, 1993). The onion cell wall contains about 42.4% pectins, 36.6% hemicelluloses and 21% cellulose (Mankarios *et al.* 1980) with 0.7% of xyloglucans and low abundance of phenolics and proteins (McCann *et al.*, 1990; McCann *et al.*, 1992).



**Figure 2.2** Schematic structural models of type I and type II walls. Cell walls illustrated are represented by Arabidopsis and rice cell walls, respectively (Yokoyama and Nishitani, 2004).

In Type I cell walls, pectic polysaccharides are one of the key components (Figure 2.3 A), which have a role in determining the pore size of the cell wall (Baron-Epei *et al.*, 1988; Carpita and Gibeaut, 1993). The pectins in plant cell walls consist of a complex set of GalA-containing polysaccharides that possess 1,4-linked  $\alpha$ -D-galactosyluronic acid (GalpA) residues (Ridley *et al.*, 2001; Caffall and Mohnen, 2009). The major types of pectins include homogalacturonan (HG), xylogalacturonan (XGA), apiogalacturonan (AGA), rhamnogalacturonan II (RG-II), and



**Figure 2.3** The formation of pectin networks by covalent and ionic bonds (Cosgrove, 2005). A shows how the pectin domains are covalently linked together to form a massively large macromolecular pectin network. This is a simplified version of a recent model by Vincken *et al.* (2003), in which rhamnogalacturonan I serves as the backbone and the other pectin domains are attached as branches. Homogalacturonans are ionically crosslinked by calcium (C) whereas boron crosslinked rhamnogalacturonan II through diester linkages (B).

rhamnogalacturonan I (RG-I) (O'Neill *et al.*, 1990; Ridley *et al.*, 2001; Caffall and Mohnen, 2009; Harholt *et al.*, 2010). Homogalacturonan (HG) is an unbranched chain of  $\alpha$ -1,4-linked-D-galacturonic acid (GalpA) residues, which is the main type of pectins in plant cell walls and the middle lamella, occupying more than 60% (Ridley *et al.*, 2001). Rhamnogalacturonan-I (RG-I) is composed of a backbone in which the disaccharide [4)- $\alpha$ -D-GalpA-(1,2)- $\alpha$ -L-Rhap] is continuously repeated (Ridley *et al.*, 2001). The backbone of GalpA residues can be substituted at various positions by  $\beta$ -D-xylosyl (Xylp) residues and  $\beta$ -D- apiofuranosyl (Apif), forming xylogalacturonan (XGA) and apiogalacturonan (AGA), respectively (Hart and Kindel, 1970; O'Neill *et al.*, 1990; O'Neill *et al.*, 2003; Zandleven *et al.*, 2006; Caffall and Mohnen, 2009). Although rhamnogalacturonan II (RG-II) and rhamnogalacturonan I (RG-I) have similar names, their structures are not related. The basic backbone of RG-II are 1,4-linked  $\alpha$ -D-GalpA residues and there are four structurally different oligosaccharide side chains linked to the RG-II backbone.

Matsuura *et al.* (2000) extracted the onion (*Allium cepa* L.) pectins and found pectins from onion contain 93.7% uronic acid, indicating the main composition of onion pectin is homogalacturonan (HG). There is also a small rhamnogalacturonan (RG) content, approximately 2.3%. Therefore, HG is very critical in onion cells. Expanding cells usually synthesize homogalacturonan which is highly methyl esterified. The degree of HG methyl-esterification plays an important role in the developmental processes of plant organs such as cell division, expansion, and adhesion (Shevell *et al.*, 1994; Wolf *et al.*, 2009). Since the un-methylated GalpA residues are anions, the un-methylated galacturonic acid residues can ionically cross-link with  $\text{Ca}^{2+}$  to form a stable gel with other pectic molecules, forming an egg-box model structure (Figure 2.3 C), which greatly increases cell wall texture and mechanical properties (Grant *et al.*, 1973; Liners *et al.*, 1989; Caffall and Mohnen, 2009; Wolf *et al.*, 2009). By contrast, although RG II molecules also can form cross-linkages, it forms covalent bonds via borate esters (Figure 2.3 B) and not ionically with calcium (Matoh and Kobayashi, 1998; Ridley *et al.*, 2001). This cross-linkage is important in the formation of normal cell walls, controlling wall porosity and wall thickness (Cosgrove, 2005). The  $\text{Ca}^{2+}$  cross-bridges condense part of the HG to create pores and contribute to the reduction in cell wall porosity (Moore and Staehelin, 1988; Knox *et al.*, 1990). The un-methylesterified HG results from the hydrolysis of the methyl ester bond of HG GalpA residues under the catalysis of pectin methylesterase (PME) secreted into the

extracellular wall space, altering the degree while pattern of methyl esterification and releasing methanol and protons (Jolie *et al.*, 2010).

### **2.2.2 Function of the Apoplast**

In general, the cell wall is a physical barrier that has several specific functions:

- (1) Maintain or determine cell shape and size to give cells a definite structure for formation of tissues and organs in plants (Geitmann, 2010). Control the rate and direction of cell growth and regulate cell volume, thereby achieving the purpose of regulating cell expansion (Bashline *et al.*, 2014).
- (2) Separate interior of the cell from the outer environment and help the cell-cell communication (Keegstra, 2010) through the plasmodesmata (specialized pores within the cell wall) (Raven, 1997) or signaling molecules located in the cell wall such as arabinogalactan proteins (Seifert and Roberts, 2007).
- (3) Constrain the turgor pressure, preventing the cell from rupturing due to turgor pressure (Knox, 2008).
- (4) Act as a physical barrier to biotic and abiotic stresses, provide protection from infection and mechanical stress, and prevent water loss (Albersheim and Anderson-Prouty, 1975; Bateman and Basham, 1976). The cell wall-degrading enzymes which are secreted by microbes can be inhibited by cell wall proteins; some enzymes within cell walls have the ability to degrade the microbes. As a result, the invasion of microorganism can be prevented by the first barrier—the cell wall (Albersheim and Anderson-Prouty, 1971; Cline and Albersheim, 1981; Laluk and Mengiste, 2010). Alterations of the structure and composition of the plant cell wall also play a crucial role in the defense mechanism against low temperature (Rajashekar and Lafta, 1996; Murai and Yoshida, 1998; Jones *et al.*, 2000; Solecka *et al.*, 2008), drought (Leucci *et al.*, 2008; Jiang *et al.*, 2012), salt stress (Uddin *et al.*, 2013; de Lima *et al.*, 2014), etc.
- (5) Provide carbohydrate storage for other metabolic processes such as seed germination (Meier and Reid, 1981; Brett and Waldron, 1996; Buckeridge *et al.*, 2000). The storage cell wall compounds in seeds include xyloglucans, pectic galactans, hemicellulose etc., which are mainly localized in the endosperm or cotyledons (Brett and Waldron, 1996). For example, under the low light conditions, the high xyloglucan content which exists in the cotyledons is used as the carbon source for the germination of *Hymenaea courbaril* (*Leguminosae*) seeds



(Santos and Buckeridge, 2004).

- (6) As economic products – many industrial products depend on cell wall material, such as timber, pulp, paper, wood, fiber, cellulosic biofuel, shelter, and even food (McNeil *et al.*, 1984; McDougall *et al.*, 1996; Bashline *et al.*, 2014).

### **2.2.3 Responses to Cold Acclimation and Role in Freezing Resistance**

As the primary physical barrier, numerous changes including the structures and compositions of cell wall are induced in response to freezing stress after a period of cold acclimation. During freezing, cell dehydration reduces cell volume but the cell wall is capable of resisting this reduction, preventing water loss and ice penetration to the intracellular spaces (Rajashekar and Burke, 1996). Dehydration resistance may lead to tightening and loosening of cell walls (Moore *et al.*, 2008b) and extensive cell wall folding as seen in *C. plantagineum* (Hartung *et al.*, 1998; Farrant, 2000) in order to avoid water loss inside cells. Water initially crystallizes in the extracellular space in plant tissues under subzero temperatures (Levitt, 1972). The plasma membrane may be damaged by the cell walls because of the mechanical distortion caused by the external ice crystals (Iljin, 1933). The ability of Jerusalem artichoke to resist freezing is enhanced through cold acclimation, although the influences are negative when the attachment between plasma membrane and cell wall is tight (Murai and Yoshida, 1998).

The apoplast can contribute to freezing resistance by physically preventing ice from nucleating the intracellular space. The quantity of cell wall increases in cold-acclimated pea seedlings (Weiser *et al.*, 1990), bromegrass cell suspension cultures (Tanino *et al.*, 1990) and leaves of winter rapeseed (Kubacka-Zebalska and Kacperska, 1999; Solecka *et al.*, 2008), contributing to cell wall thickening to a certain degree (Solecka *et al.*, 2008). Reports of cell wall thickening during cold acclimation are consistent with the idea that altered cell wall elasticity is considered an important component of the adaptive response (Hüner *et al.*, 1981), which is thought to affect water movement and in terms of freezing resistance, related to the pattern of ice propagation in the cold-acclimated plant tissues (Griffith and Brown, 1982; Griffith *et al.*, 1985). Cell wall thickening is also observed in the leaves of potato (Chen *et al.*, 1977), cold-hardy rye cv. Puma (Huner *et al.*, 1981; Griffith and Brown, 1982) and winter oilseed rape (Stefanowska *et al.*, 1999) in response to cold acclimation. Cell wall strength increases and the pore-size of the cell wall decreases in grape and apple cells upon cold acclimation (Rajashekar and Lafta, 1996).

Increased cell wall rigidity has been proposed to protect cells against intracellular freezing (Rajashekar and Lafta, 1996).

Alterations in cell wall composition, the activities of cell wall-modifying enzymes, and the accumulation of particular proteins in the apoplast space in plant tissues are induced during cold acclimation (Weiser *et al.*, 1990; Marentes *et al.* 1993, Hon *et al.* 1994, Antikainen *et al.* 1996). The water soluble carbohydrate compositions including fructan, sucrose and fructose of cereal crops also changes during exposure to freezing temperatures, in which fructan decreased while fructose and sucrose increased (Olien and Lester, 1985; Olien and Clark, 1993). As a result of fructan hydrolysis, the increase in sugars in the apoplast is thought to be one of the reasons for the improved hardness during subzero acclimation (Olien and Lester, 1985; Livingston and Henson, 1998). The presence of cell wall polymers reduces the growing speed of ice crystals in the middle lamella (Olien and Smith, 1977). Studies have demonstrated that the epoxide monomer, lipid-derived polymers cutin and suberin are deposited and can be visualized in the mestome sheath cell walls (Kolattukudy, 1980). Moreover, cold acclimation increases hemicellulose content (Weiser, 1990; Kubacka-Zębalska and Kacperska, 1999) and reduces xylose and glucose contents in winter oilseed rape leaves (Kubacka-Zębalska and Kacperska, 1999). Cold acclimation also affects the accumulation of cell-wall proteins and the activities of cell-wall enzymes. Extensin, one of the structural cell wall glycoproteins which are thought to provide rigidity to the cell wall by forming an interpeptide-linked network (Lampert, 1967 ), increases in cold acclimated pea along with an increase in freezing resistance (Weiser *et al.*, 1990). Two glycine-rich proteins encoded by *MsaCiA* and *MsaCiB* mRNAs accumulate in both leaf and crown cell walls of cold acclimated alfalfa (Ferullo *et al.*, 1997). Numerous other studies have focused on antifreeze proteins (AFPs) which accumulate in the mesophyll cell walls, secondary cell walls of xylem vessels and epidermal cell walls of plants during cold acclimation (Griffith *et al.*, 1997). Cold acclimation increases both transcriptional and translational level of AFP genes (Yeh *et al.*, 2000; Huang and Duman, 2002). AFPs can adsorb onto ice crystals and prevent ice growth (Raymond and DeVries, 1977). AFP activity is present only in the apoplastic extracts from freezing tolerant monocotyledons after cold acclimation (Antikainen and Griffith, 1997), where they enhance freezing resisting (Griffith and Yaish, 2004; Griffith *et al.*, 2005). Apoplastic AFPs modifies the normal growth pattern of ice crystals in winter rye after cold acclimation and endures extracellular ice during freezing stress (Griffith *et al.*, 1992; Hon *et al.*,

1995). The study by Griffith *et al.* (1992) illustrates that proteins with antifreeze-like properties from the apoplast of acclimated rye leaves have antifreeze activity and these protein mixtures modify the growth pattern of ice crystals. Interestingly, AFPs do not exhibit antifreeze activity in non-acclimated plants even though they are present within those plants (Wang *et al.*, 2006). The antifreeze activity of AFPs is activated after exposure to low temperature and is only present in plants which can tolerate the presence of ice in their tissues (Yaish *et al.* 2006). Some cell-wall degrading enzymes are also modified by cold acclimation. For instance, the activity of pectin methylesterases (PMEs) was enhanced during cold acclimation in winter oilseed rape (Solecka *et al.*, 2008) and in cold-treated *Arabidopsis* (Lee, 2003). This enzyme is capable of regulating the degree of pectin methyl esterification through catalysing the demethylation reaction of methylesterified pectin to form the un-methylesterified pectin, affecting the cell wall rigidity.

These observed changes in cell wall compositions are associated with the regulation of genes encoding proteins involved in cell wall modification (Zabotin *et al.*, 1998). The same genes may be regulated in various ways at low temperature. For example, the expression of the *EXP* genes (expansin genes which modulate cell-wall loosening and their activities can be affected by drought (Jones and McQueen-Mason, 2004) and salt stress (Buchanan *et al.*, 2005)). During cold acclimation, *EXP* genes are down-regulated in *Arabidopsis* roots (Swindell, 2006) and sweetpotato (Noh *et al.*, 2009), but up-regulated in *Arabidopsis* seeds and rice anthers (Imin *et al.*, 2004; Yamauchi *et al.*, 2004). Both *EXP* and *XTH* (xyloglucan endotransglucosylase / hydrolase genes), which regulate cell wall xyloglucan content and plant growth (Miedes *et al.*, 2013), are known to improve resistance to drought, salt and aluminum stresses (Han *et al.*, 2013). *EXP* and *XTH* genes in sweetpotato, Extensin and *XTH* encoding genes in *Arabidopsis* are down-regulated under cold response, consistent with the improvement in freezing tolerance (Seki *et al.*, 2002). Pectin methylesterases (PMEs) located in the cell walls are encoded by a multigene family (Richard *et al.*, 1996; Micheli *et al.*, 1998), and 67 *PME*-related genes have been identified from the *Arabidopsis* genome (The *Arabidopsis* Genome Initiative, 2000). *PME* gene expression or its activity is generally thought to be strongly associated with dormancy breakage (Ren and Kermode, 2000), seed germination (Müller *et al.*, 2013), tube pollen growth (Wakeley *et al.*, 1998; Futamura *et al.*, 2000), and fruit maturation (Kagan-Zur *et al.*, 1995). In addition, Zhu *et al.* (2013) demonstrated a *PME3* in cold acclimated cotton is up-regulated. Arabinogalactan proteins (AGPs) located in plant cell walls, which are involved in plant growth

and development, are also up-regulated in response to cold stress during early root development of cotton (Gong *et al.*, 2011). Although numerous researchers support cell walls functions as a barrier and provides freezing resistance, relatively little work has been done to identify those specific components within the cell wall and the apoplastic space that are critical to this resistance.

Pectic polysaccharides are connected to the integrity and rigidity of plant tissue, cell wall porosity (Baron-Epel *et al.*, 1988) and abiotic and biotic stress resistance (Hahn *et al.*, 1989; Vorwerk *et al.*, 2004), affecting the tissue ability to supercool (Ashworth and Abeles, 1984). In the study of Solecka *et al.* (2008), the tensile strength of a leaf tissue was positively associated with pectin content and its degree of methyl esterification after cold acclimation. Cell wall porosity and gelation is generated via the formation of the pectic gel resulting from ionically cross-linking un-methylesterified HG and  $\text{Ca}^{2+}$  and the degree of pectin methyl-esterification (Grant *et al.*, 1973; Yoo *et al.*, 2003). Cell wall porosity is related to the resistance of plant cells to low temperature damage. For example, the pore size of the cell wall in both grape and apple cell cultures decreased and the cell wall strength increased after cold acclimation (Rajashekar and Lafta, 1996). The ultrastructural observation of peach and dogwood cells demonstrated that both pit membrane (black cap and primary cell-wall) and xylem parenchyma cells were almost completely digested by the treatment of macerage (an enzyme mixture rich in pectinase), resulting in a complete shift of the low-temperature exotherm (LTE) to warmer temperature as determined by differential thermal analysis (DTA) (Wisniewski *et al.*, 1991a). Thus, pectins appear to regulate water and ice movement through vascular pits thereby enhancing deep supercooling (Wisniewski *et al.*, 1991a). This result was further confirmed by the study from Jones *et al.* (2000) in which a 3000 MW dextran-conjugated dye cannot penetrate the bud axes after cold-acclimation, indicating that this lower porosity probably restricts the penetration of ice into buds from the grape cane. In addition, Jones *et al.* (2000) found apoplastic permeability increased because of the treatment of pectinase which removes pectin, and increased the ice nucleation temperature, preventing supercooling according to the results from DTA. Using FTIR, Wilson and his co-workers (2000) indicated macromolecules in onion cell walls such as cellulose and pectin are reoriented along the direction of cell elongation (parallel) from a small directional alignment with parallel and perpendicular polarizations after going through a mechanical stress.

The pectin content increases in pea epicotyls (Weiser, 1990), winter oilseed rape leaves (Kubacka-Zębalska and Kacperska, 1999; Solecka *et al.*, 2008) and bromeliad (*Nidularium minutum*) (Carvalho *et al.*, 2013) during cold acclimation. In the frost-tolerant pea genotype, a period of cold acclimation induced an increase in HG (Baldwin *et al.*, 2014). The degree of pectin methylesterification is regulated by the pectin methylesterases (PMEs). PME activity increases in cold-acclimated oilseed rape leaves, which was associated with the improved cell wall rigidity caused by de-methylesterification of pectin. More un-methylesterified pectin was formed as a result of higher PME activity, creating a semi-rigid pectate gel through the  $\text{Ca}^{2+}$  cross-linking with free carboxyl groups (Jarvis, 1984; Carpita and Gibeaut, 1993; Solecka *et al.*, 2008). Therefore, the enhanced PME activity could improve cell wall stiffness and increase freezing tolerance of plants. In addition, as ‘pectic plasticizers’, the arabinose polymers can hold the flexibility of cell walls during dehydration and rehydration of guard cells (Jones *et al.*, 2003) and aquatic plants (Moore *et al.*, 2008a) and thus may protect cells from damage in the process of freezing-thawing.

## **2.3 Calcium**

### **2.3.1 Calcium is an Essential Element in Plants**

Calcium is one of 14 essential nutrients in plants and plays important roles in structural support of the cell wall and plasma membrane, growth and signal transduction in plant tissues (Maathuis, 2009). As the cellular structural supporter, the cation  $\text{Ca}^{2+}$  can form the complexes with anion groups of organic compounds including phosphates and carboxyl of phospholipids, proteins and sugars. In the process of cell wall formation,  $\text{Ca}^{2+}$  is a crucial element to determine the structural rigidity of the cell wall since  $\text{Ca}^{2+}$  can electrostatically form cross linkages to the free carboxyl groups of acidic pectin residues which are released by hydrolysis of the methyl-esterified pectin under the catalysis of PME (Wyn Jones and Lunt, 1967; Burström, 1968; Hepler, 2005). Calcium in plant cells mainly localizes in the endoplasmic reticulum (about 1mM) and the apoplast (around 1-10mM) (Bush, 1995; Trewavas and Malho, 1998). The high apoplastic  $\text{Ca}^{2+}$  concentration, up to 60-75% of the total tissue  $\text{Ca}^{2+}$  content (Demarty *et al.*, 1984), makes the cell wall more rigid and less plastic, helping to increase biotic and abiotic stress resistance (Hepler, 2005). In the plant plasma membrane,  $\text{Ca}^{2+}$  complexes with phosphate groups from

phospholipids in order to maintain the integrity and permeability of the plasma membrane (Hepler, 2005; Maathuis, 2009).

Calcium is necessary at all stages of plant growth and development including the regulation of growth and differentiation in cells and tissues (Hepler and Wayne, 1985; Trewavas and Malho, 1998; Sanders *et al.*, 1999), embryogenesis (Medvedev, 1997) etc.  $\text{Ca}^{2+}$  also acts as a secondary messenger. Changes to the cytosolic free  $\text{Ca}^{2+}$  in plants are induced by the biotic and abiotic stress, stomatal regulation and physical damage (Mahouachi *et al.*, 2006; McAinsh and Pittman, 2009). Research has demonstrated that  $\text{Ca}^{2+}$  influx is temporally moved into the cell cytoplasm under the stimuli of the stress signal, inducing a transient increase in cytosolic  $\text{Ca}^{2+}$  which then activates the cascades of phosphoproteins like calmodulin and finally induces the expression of stress-responsive genes such as antifreeze proteins (AFPs) during freezing stress (Sanders *et al.*, 1999; Knight, 2000, Xiong *et al.*, 2002).

### **2.3.2 Role of Calcium in the Apoplast and Freezing Stress Resistance**

Today, no one questions that  $\text{Ca}^{2+}$  is a crucial factor in regulating the growth and development of plants. Through coupling of extracellular signals and intracellular second messengers,  $\text{Ca}^{2+}$  participates in various physiological and biochemical reactions such as controlling plant growth and differentiation (Hepler and Wayne, 1985), regulating embryogenesis (Medvedev, 1997), assembling and disassembling of cytoskeleton elements (Medvedev and Markova, 1998), and modulating movement of stomatal cells (Ng *et al.*, 2001). Thus,  $\text{Ca}^{2+}$  plays vital roles in the whole life cycle of plant (Medvedev, 2005). Intracellular levels of calcium ions are significantly modulated following a plant's exposure to either abiotic or biotic stresses (Ludwig *et al.*, 2004). For example, cold stimuli can induce a rapid increase in plant cellular  $\text{Ca}^{2+}$  level (Knight *et al.*, 1996). The cold resistance in grafting eggplant seedlings after the application of exogenous  $\text{Ca}^{2+}$  is higher than those with calcium deficiency (Gao *et al.*, 2004).

Since the 19th century,  $\text{Ca}^{2+}$  has been known to play a crucial role in determining the structural rigidity and stabilization of the cell wall (Wyn Jones and Lunt, 1967; Burström, 1968) through forming calcium bridges with pectin molecules which is regulated by the activity of PME (Chang *et al.*, 1995). Suutarinen *et al.* (2000) used instrumental textural measurements, as well as microscopic studies to show that in most cases,  $\text{CaCl}_2$  pre-freezing treatments stabilized the structures including cell walls (in particular pectin) of strawberry tissues. Two areas within

the cell have been recognized as being important targets for  $\text{Ca}^{2+}$ : the cell wall and endoplasmic reticulum. In terms of the cell wall, calcium is a major pectin cross-linking agent which results in increased gel density, reduced cell wall porosity and increased cell wall rigidity (Peaucelle *et al.*, 2012). Therefore, we propose that the dynamic increase in pectin gelation which is driven by PME activity and calcium binding during cold acclimation provides cells ability to avoid freezing damage through a physical blockage of ice crystals from nucleating the intracellular space.

## **2.4 *Allium fistulosum* L.**

*Allium fistulosum* L. ( $2n = 2x = 16$ ) (Inden and Asahira, 1990), known as Japanese bunching onion or Welsh onion, is a perennial plant and belongs to the genus *Allium* L. in the section *Cepa* (Mill.) Prokh (Hanelt, 1990) and is one of the most important cultivated edible *Allium* crops (Others are onion, leek, garlic.). Other names that may be applied to this plant include green onion, spring onion, scallion, and salad onion. It is thought to have originated in northwestern China and is a leaf vegetable mainly cultivated in East Asian countries, such as China, Japan and Korea, as well as in European countries (Grevsen, 1989; Ford-Lloyd and Armstrong, 1993). The Japanese onion contains abundant vitamin C (31-36%) (Kołota *et al.*, 2013), but also flavonoids (283 mg/kg Fresh Weight) (Aoyama and Yamamoto, 2007), macro- and micro-nutrients, especially Ca (1%) and K (3%) (Kołota *et al.*, 2013). Most onions can multiply by forming perennial evergreen clumps (Thompson, 1995).

Japanese bunching onion is vigorous with a rapid growth rate (Lazić *et al.*, 2002; Štajner *et al.*, 2006) and resists summer drought, winter cold and poor soil quality (Yamasaki *et al.*, 2003). Some Japanese bunching onions can survive when the air temperature drops below  $-40^{\circ}\text{C}$ . The species also has the resistance ability against onion blight, pink root (Netzer *et al.*, 1985), and other plant diseases and insect pests. Onion has a unicellular epidermal layer (Wilson *et al.*, 2000) located at the base of the leaf and is easy to peel. In addition, the individual cell size of Japanese bunching onion is quite large, approximately  $250\text{ }\mu\text{m}$  in length x  $50\text{ }\mu\text{m}$  in width x  $90\text{ }\mu\text{m}$  in thickness. The epidermal cell layer has no chlorophyll, avoiding issues of auto-fluorescence during cell studies based on fluorescent probes. Therefore, *Allium fistulosum* (Japanese onion bunching) is an ideal system to observe freezing processes.

In 1956, Asahina (1956) conducted freezing tests on the epidermis of unhardy and hardy Welsh onion leaves. He found that the unhardy cells became immediately opaque and dark and then lighter once the temperature decreased to below the freezing temperature, and the ice would rapidly propagated the whole protoplast. In contrast, the freezing rate in epidermal cells was reduced and the darkening phenomenon in frozen cells was lowered if the epidermal cells are hardy or have a certain degree of desiccation. In those studies, intracellular freezing (at  $-13.5^{\circ}\text{C}$ ) in hardy epidermis of Welsh onion was markedly later than extracellular freezing (at  $-7^{\circ}\text{C}$ ). These interesting observations led us to study the mechanism of intracellular freezing stress avoidance on *Allium fistulosum*.



### 3.0 MATERIALS AND METHODS


#### 3.1 Plant Materials and Growth

Japanese bunching onion (*Allium fistulosum* L.) seeds produced from plants growing outdoors in Saskatoon, SK (52°7'N) were collected and germinated in the College of Agriculture and Bioresources greenhouses (45 Innovation Blvd., University of Saskatchewan, Saskatoon, SK Canada). Plants were grown for 3 months in 6" pots of soilless mix (Sunshine No. 4, Sungro Hort Inc, Bellvue, WA) at (20 ± 5°C) under natural light supplemented with 400 W high-pressure sodium lights (18 h photoperiod, average of 600  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ ) (see Appendix A). Plants were irrigated once every second day and fertilized weekly with 20-20-20 (NPK, 2 g·L<sup>-1</sup>) + micronutrients including boron, copper, iron, manganese, molybdenum, zinc (Plant Products Co. Ltd., 314 Orinda Road, Brampton, Ontario, LGT-1J1). For calcium treatment, each plant was watered with 100 ml 0.05 M CaCl<sub>2</sub> solution every second day over a four-week periods in the greenhouse. Subsequently, half of the treatments plus and minus calcium were transferred into a Conviron growth chamber with 12°C/4°C (day/night) under an 8 hour photoperiod with light intensity of 370  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  for two weeks for cold acclimation treatment. In all, there were four different treatments: Ca (-) non-acclimated (NA), Ca (+) non-acclimated (NA+Ca), Ca (-) acclimated (ACC), Ca (+) acclimated (ACC+Ca).


#### 3.2 Freezing Stress Treatments

The leaves of onion were numbered from 0 to 4 in order of age in which 0 is the youngest leaf and 4 is the oldest leaf (Figure 3.1, also see Appendix I). In grass, the collar region is located between the leaf blade and leaf sheath, containing structures called ligules which is the membranous tissue located at the junction of the leaf blade and leaf sheath (Figure 3.2) (Harper and Freeling, 1996). The ligule is also called epidermal layer which is easily peeled off from onion and kept as an intact single cell layer, simplifying sample preparation. Bin Li previously performed the freezing test among different ages of leaves and different regions of the same ligule (Figure 3.1). More specifically, the freezing tolerance in the leaves was found to be leaf0 > leaf1 > leaf2 > leaf3 > leaf4. The region C-b located between 2 and 3 cm from the ligule in the center of the sheath epidermal cell layer (Figure 3.1 B) is more freezing stress resistant than other regions of the same epidermal layer (Tanino *et al.*, 2013). However, the region B-b (located

between 1 and 2 cm from the ligule in the center of the sheath epidermal cell layer) in leaf2 located in the center of the various regions expressed greater consistency of freezing stress resistance and was selected in this study. Leaf 2 of each sample (NA, NA+Ca, ACC and ACC+Ca) was placed into a 50 ml glass tube with 2 ml distilled water, and cooled to  $-2.5^{\circ}\text{C}$  for 2 hours using an Endocal RTE-Series Refrigerated Bath. Thereafter, ice was added to the tube to nucleate the sample. After one hour, the temperature was decreased at a rate of  $-5^{\circ}\text{C}\cdot\text{hour}^{-1}$ , held for one hour and the first sample was removed. Samples were removed at  $-10^{\circ}\text{C}$ ,  $-15^{\circ}\text{C}$ ,  $-20^{\circ}\text{C}$ , and  $-25^{\circ}\text{C}$ ,  $-30^{\circ}\text{C}$ . Upon removal from the refrigerated bath, the samples were placed into a  $4^{\circ}\text{C}$  refrigerator overnight to thaw. The next morning, samples were warmed to room temperature ( $24^{\circ}\text{C}$ ) for at least one hour before observation. The epidermal layer of each sample was peeled and fixed onto the slide with a coverslip. Presence or absence of protoplasmic streaming was used as a measurement of the number of live and dead cells (10 cells per sample, 3 samples per replication, and 5 replications per temperature).



**Figure 3.1** The experimental area of Japanese bunching onion. A: numbering system of the whole plant according to leaf age; B: numbering system in one ligule; C: the ligule (or epidermal layer) separated from the sheath; D: the ligule epidermal layer placed on a glass slide.



**Figure 3.2** The ligule and collar region of the plant. Right: the classical structures of ligule in Poaceae (<http://weeds.cropsci.illinois.edu/weedid.htm>). Left: the ligule in *A. fistulosum*.

### 3.3 Analysis of Freezing Process

The freezing process including ice nucleation in extra- and intra-cellular space was recorded with a Nikon E400 microscope in Department of Food and Bioproduct Sciences, University of Saskatchewan. The cryostage apparatus (Linkam freezing stage, LTS 120, UK) was connected to a Nikon E400 light microscope with a camera attached (Nikon Digital Sight). The epidermal layers of leaf 2 from control and treated onions were peeled and the abaxial surface placed down onto the glass slides. The slide was fixed on the cryostage. Before lowering the temperature, protoplasmic streaming of live cells for viability analysis was confirmed and video recorded at 24°C. The cryostage was water-cooled using a heat sink pump and the cooling rate was adjusted by a controller (Linkam PE94, UK). The temperature was decreased to 0°C and finally to -23°C (the lowest temperature achieved by the heat sink) at a rate of 5°C·hour<sup>-1</sup>. The process of ice crystal formation in the cells was video-recorded in real time with Cam Studio using a Nikon 40X lens (see Appendix B). When ice formation occurred, the refraction of light changed and the temperature at which extra- and intra-cellular ice nucleation occurred was recorded. Samples were then thawed at a rate of 1°C·min<sup>-1</sup> and again examined for the presence of protoplasmic streaming (see Appendix C) when temperature reached 24°C. Treatments also included ice nucleation (+) induction by adding a few grains of silver iodide powder (AgI) resuspended in water to the glass slide. In these experiments, the onion epidermal layer was placed on top of the silver iodide. The temperature was then decreased to -2.5°C and held for two hours to further initiate ice nucleation. Each treatment was replicated five times.

### 3.4 Study of Apoplast Permeability

The epidermal layers of non-acclimated (NA) and acclimated (ACC) onions were

submerged in the mixture of fluorescein conjugated to a 3,000 MW and 70,000 MW dextran solution ( $5 \text{ mg} \cdot \text{ml}^{-1}$  in 0.1 M phosphate buffer, Life Technologies) for 16 h and observed by a Laser Scanning Confocal Microscope (ZEISS LSM 510 Confor2, Department of Biology, University of Saskatchewan, see Appendix E).

The diffusion rate of fluorescein (Sigma-Aldrich Chemical Co. St. Louis, MO) from the extracellular to the intracellular space was measured using a Laser Scanning Confocal microscope (LSM, Nikon Eclipse 80i Microscope, see Appendix E). The epidermal layers of non-acclimated and cold acclimated onions were peeled and flattened onto the cover well with glass wool (the adaxial side was toward the glass wool), and then covered with a glass slide. After sample preparation, the slide was placed into the LSM. Next, the  $100 \text{ } \mu\text{l}$   $1 \text{ mg} \cdot \text{ml}^{-1}$  fluorescein solution was slowly injected through a hole in the cover well to soak. The focus was kept approximately  $40 \text{ } \mu\text{m}$  which roughly corresponds to the midpoint of the intracellular space (total onion cell thickness is  $90 \text{ } \mu\text{m}$ ). The diffusion process of fluorescein in individual samples were recorded every 10 seconds by the NIS-Elements imaging software. In order to calculate the diffusion rate of the samples, the NIS-Elements software was used to select the region of interest (ROI) with an intact cell including the apoplastic space, plasma membrane and cytoplasmic region. The ROI in the samples was scanned and the time at which the pixel brightness increased to 2000 pixels (mean intensity) at that location was recorded. The diffusion rate was calculated based on the slope with pixels as the Y axis and time as the X axis.

### **3.5 Analysis of Pectin Methyl Esterification**

#### **3.5.1 Sample Preparation and Fixation**

The sample preparation was based on Lee *et al.* (2008). The onion whole sheath was cut into 2-3 mm pieces and fixed in 0.01 M sodium phosphate buffer (including 1% formaldehyde and 0.025% glutaraldehyde, pH 7.2). The samples were infiltrated with vacuum for a minimum of 1 h in order to thoroughly fix the samples and then incubated at  $4^{\circ}\text{C}$  overnight. The fixed samples were then successively washed with 0.01 M sodium phosphate buffer twice for a total of 4 hours and dehydrated in a graded ethanol series (30, 50, 70, 95, and 100%) at one hour for every concentration of ethanol at  $4^{\circ}\text{C}$ . The samples were then infiltrated with London Resin (LR) white (Electron Microscopy Sciences, <https://www.emsdiasum.com>):ethanol (1:1) for 5 hours at  $4^{\circ}\text{C}$ , and this step was repeated once overnight. The samples were subsequently transferred into a

2:1 concentration of LR white resin: ethanol for 7 hours and then overnight. The last infiltration step with pure LR white resin was added to the samples at least 6 times and changed every 12 hours. Finally, the samples were transferred into an embedding mold and polymerized at 60°C for 24 to 48 hours. The samples were sectioned after embedding using the Reichert-Jung Ultracut Microtome equipped with a diamond knife (Department of Biology, University of Saskatchewan). Sections (1 µm thick) were placed on poly-L-lysine coated slides (Electron Microscopy Sciences, <https://www.emsdiasum.com>) and heated on a 55°C warm plate to adhere the sections to the slide.

### **3.5.2 Analysis of Pectin Methyl Esterification by Immunohistochemistry**

The immunofluorescence labelling analysis to evaluate the degree of pectin methyl esterification was conducted with the antibodies JIM5 recognizing de-methylesterified epitopes of HG and JIM7 recognising methylesterified epitopes of HG (CarboSource Services, [http://www.ccrcc.uga.edu/~carbosource/CSS\\_home.html](http://www.ccrcc.uga.edu/~carbosource/CSS_home.html)). These experiments were conducted in the Norwegian University of Life Sciences Imaging Lab, Ås, Norway and subsequently repeated three times in the University of Saskatchewan. The experimental procedure followed Lee *et al.* (2008). The 1 µm thick sections obtained as described above were processed and were first incubated in 1 ml 3% (w/v) milk protein in phosphate-buffered saline (PBS), pH 7.2, for at least 30 min to block non-specific binding sites. Thereafter, sections were incubated with primary rat monoclonal antibodies JIM5 and JIM7 diluted 1:10 in PBS, respectively, for at least 1 h at room temperature and then washed three times with PBS at least 5 min for each time. Next, the secondary anti-rat-IgG+IgM+IgA antibodies linked to fluorescein isothiocyanate (FITC; Abcam) were diluted 100-fold PBS and 200 µl was added to each slide. After 1 h incubation at room temperature, the sections were washed in PBS buffer three times at least 5 min for each time. Finally, the sections were mounted with prolonged gold antifade reagent (Life Technologies) and examined with a fluorescent microscope (Evos digital inverted microscope, see Appendix E). All incubation steps were operated in the dark. The blank control was only added to the secondary antibody in order to show that the label is specific to the primary antibody and the other steps followed above.

### **3.5.3 Determination of PME Activity**

PME activity under four different treatments was performed according to Richard *et al.*

(1994). Fresh epidermal layers (0.2 g) of leaf 2, the non-epidermal layer (2 g) of leaf 2 and the whole sheath (including both epidermal layer and non-epidermal layer (2.2 g) of leaf 2 (see Appendix A) were homogenized in 0.05 M HEPES buffer, pH 6.5, containing 1X protease inhibitor cocktail (P9599, Sigma-Aldrich Chemical Co. St. Louis, MO), and washed several times with cold water. After air drying, crude cell wall preparation was used to extract cell wall proteins with 0.05 M HEPES buffer containing 1 M NaCl and 1X protease inhibitor cocktail, under gentle agitation at 4°C for 2 h. Protein concentration in the extracts was determined by the method of Bradford (1976) using a spectrophotometer (GENESYS 10 Bio, Thermo Fisher Scientific, USA) and using different concentrations of bovine serum albumin (Sigma-Aldrich Chemical Co. St. Louis, MO) as standards. The prepared protein extracts from cell walls were used to determine PME activity.

In order to measure the PME activity, the reaction mixtures contained 0.5% (w/w) highly methylated citrus pectins (Sigma Chemical Co. St. Louis, MO), 0.2 M NaCl and 0.015 (w/v) methyl red as a pH indicator. The extract above (5 µl) was added to 1 ml of reaction mixture. Changes in colour from yellow to red (due to reduced pH during pectin de-esterification) were measured spectrophotometrically at 525 nm for 2 min at 25°C. A calibration curve was generated by adding 1-200 nEq H<sup>+</sup> from HCl to 1 ml of reaction mixture and the enzyme activity was calculated on the basis of cell wall protein. PME activity was expressed in PME units where one unit was defined as one nanoequivalent of protons (nEq H<sup>+</sup>) released by milligrams of cell wall proteins during 1 min and recorded as a function of the original FW of that tissue.

The PME activity in the cooking onion (*A. cepa*) and green onion (*A. chinense*) were compared with Japanese bunching onion (*A. fistulosum*) and determined as described above.

### **3.6 Cell Wall Compositional Analysis**

The cell wall compositional analysis followed the methods of Foster *et al.* (2010). These measurements were conducted in conjunction with RWTH Aachen University, Germany.

#### **3.6.1 Cell wall isolation**

The epidermal layers of non-acclimated and acclimated onions were peeled and freeze dried. Roughly 15 mg of sample was ground with 5.5 mm stainless steel balls in a 2 ml sarstedt screw cap tube for 1 min using a Retsch mill (Retsch GmbH, Haan, Germany) operating at 25 Hz. After removing the steel balls, the ground samples were vortexed thoroughly with 1.5 ml 70%

aqueous ethanol and centrifuged at 10,000 rpm for 10 min to pellet the alcohol insoluble residue. The supernatant was decanted and 1.5 ml chloroform/methanol (1:1 v/v) solution was added to re-suspend the pellet by shaking the tube thoroughly. The mixture was then centrifuged at 10,000 rpm for 10 min and the pellet was re-suspended in 500  $\mu$ l acetone. The solvent was evaporated to dryness at 35°C. In order to remove the starch from the cell wall, the pellet was re-suspended in 1.5 ml 0.1 M sodium acetate buffer (pH 5.0) and heated for 20 min at 80°C and then cooled on the ice. Next, the pellet was added with agents including 35  $\mu$ l 0.01% sodium azide ( $\text{NaN}_3$ ), 35  $\mu$ l  $\alpha$ -amylase (50  $\mu$ g/1ml  $\text{H}_2\text{O}$ ; from *Bacillus licheniformis*, Sigma-Aldrich Chemical Co. St. Louis, MO) and 17  $\mu$ l pullulanase (1.25 g/ml, 18.7 units from *Bacillus acidopullulyticus*, Sigma-Aldrich Chemical Co. Inc. St. Louis, MO), and incubated overnight at 37°C in the shaker. The suspension was heated to 100°C for 10 min to terminate digestion before centrifuging at 10,000 rpm for 10 min. The supernatant containing solubilized starch was discarded. Meanwhile, the pellet was washed with 1.5 ml water three times and the pellet was re-suspended in 500  $\mu$ l acetone. The prepared cell wall material was pelleted after air drying.

### **3.6.2 Matrix Polysaccharide composition**

To determine the cell wall monosaccharide composition, 2 mg cell wall material (obtained as described above) was placed into 2 ml starstedt tubes and 20  $\mu$ l inositol solution (5 mg/ml) was added as an internal standard. The cell walls were rinsed with 250  $\mu$ l acetone to collect the cell wall material at the bottom of the tube and the acetone was evaporated very gently under airflow.

For the weak acid hydrolysis, each sample was mixed with 250  $\mu$ l 2 M trifluoroacetic acid (TFA) and incubated for 90 min at 121°C. The sample was then cooled on ice and centrifuged at 10,000 rpm for 10 min. About 100  $\mu$ l acidic supernatant, which contained the matrix polysaccharide derived monosaccharides, was transferred to 2 ml glass screw cap vials and the remaining pellet was used for the crystalline cellulose assay. The TFA was evaporated from the glass tube under gentle air. Then 300  $\mu$ l 2-propanol was added, vortexed and evaporated at 25 °C. This step was repeated three times. Each dried sample had 200  $\mu$ l of a sodium borohydride solution added since the reduction of the monosaccharides is the first step of the alditol acetate derivatization process. The fresh solution with 10 mg sodium borohydride per 1 ml of 1 M ammonium hydroxide was prepared every time and the glass vial was placed at room temperature for 1.5 hours. The solution was neutralized with 150  $\mu$ l of glacial acetic acid before


vortexing and evaporating at 25°C. Then, 250 µl acetic acid/methanol (1:9 v/v) was added. After vortexing and evaporating at 25°C, 250 µl methanol was added and evaporated, and this process was repeated three times. Two chemicals including 50 µl acetic anhydride and 50 µl pyridine were added to the glass vial and incubated at 121°C for 20 min in order to make the acetylation of the alditols. The samples were cooled down to room temperature and the reagents were then evaporated from the samples by air flow. The next step of adding 200 µl toluene to evaporate, was repeated three times. The final steps extracted the alditol acetates. First, 500 µl ethyl acetate was added and slightly swirled. Second, 2 ml water was added and vortex very well. To obtain the clear separate layers, the mixture was centrifuged at 2,000 rpm for 5 min. Third, 50 µl of the ethyl acetate layer was diluted with 100 µl acetone and pipetted into GC/MS vials with inserts. Finally, the samples were injected into a GC equipped with a quadrupole MS (Agilent Technologies, Santa Clara, United States). A Supelco SP-2380 (30 x 0.25 mm x 0.25 µm film thickness) column was used with a 4 min solvent delay and a flow rate of 1.5 ml/min. Injected samples were subjected to the following temperature program: initial hold at 160°C for 2 min; a 20°C/min ramp to 200°C and hold for 5 min; a 20°C/min ramp to 245°C and held for 12 min; spike to 270°C and hold for 5 min before cooling to the initial temperature of 160°C. Peaks were identified by mass profiles and/or retention times of standards including fucose, rhamnose, arabinose, galactose, glucose, xylose, mannose, galacturonic acid and glucuronic acid. Monosaccharides were quantified based on standard curves.



## 4.0 RESULTS AND DISCUSSION

### 4.1 Freezing Stress Resistance


Japanese bunching onions were acclimated in the growth chamber with 12/4 °C (day/night) for one- and two-weeks. The acclimation temperatures used in this project simulated the natural temperature changes in autumn in Saskatoon, SK Canada. In addition, the pre-test showed the leaves of Japanese bunching onion turned yellow under 8/4 °C (day/night) after four weeks acclimation, because onions in nature survive in winter through leaf dry down to re-mobilize carbohydrates to store in the bulb. Freezing resistance in *Allium fistulosum* epidermal cell layer increased under cold acclimation (ACC) (Figure 4.1). Presence of protoplasmic streaming was used as a viability test for live cells, whereas absence of protoplasmic streaming was interpreted to indicate cell death. After two weeks of cold acclimation under 12/4 °C (day/night), the LT<sub>50</sub> (lethal temperature at which 50% of the cells were killed) of the epidermal cell layer of Japanese bunching onions decreased to -27°C, compared to non-acclimated (NA) onions which had an LT<sub>50</sub> value of -12°C (Figure 4.1). This cold acclimation response was confirmed independently in Japan using the same source of *Allium fistulosum* but acclimated at a constant 2°C for two weeks (Kobayashi *et al.*, 2014). In the Japanese study, the vital stain, BCECF (2',7'-Bis-(2-Carboxyethyl)-5-(and-6)-Carboxyfluorescein) was used to determine viability. BCECF is a membrane-permeable fluorescent indicator which can be converted into fluorescent products (Kobayashi *et al.*, 2014). These products can either permeate into the cell freely by intracellular esterases or be retained by cells due to their negative charges ([https://www.dojindo.com/Protocol/CellStaining\\_Protocol.pdf](https://www.dojindo.com/Protocol/CellStaining_Protocol.pdf)). Freezing stress resistance in this Japanese study also corresponds to our earlier work in which the *Allium fistulosum* epidermal cell layer was determined to be alive at -25°C after two weeks of cold acclimation (Tanino *et al.*, 2013). Therefore, using *Allium fistulosum* epidermal cell layers as a model for cold acclimation and freezing stress resistance appears to be reliable and consistent.



**Figure 4.1** Effect of cold acclimation time on freezing resistance in non-acclimated (NA), 1-week and 2-week acclimated epidermal layer of *Allium fistulosum* L. (ACC). Cold acclimation was done using 12/4°C, 8 hr photoperiod. Error bars are means  $\pm$  SD based on the five separate experiments.

After the application of 0.05 M  $\text{CaCl}_2$  solution as a soil drench for two weeks, the  $\text{LT}_{50}$  in non-acclimated onion cells (NA+Ca) was about  $-14^\circ\text{C}$ . As compared to no (minus) calcium treatment (NA) ( $-12^\circ\text{C}$ ), and there was no significant difference (Figure 4.2). However, some NA+Ca cells were still alive at  $-20^\circ\text{C}$  in non-acclimated onions. In the absence of cold, a few reports have shown that the application of calcium alone can increase the freezing resistance in plants such as evergreen oak and apple (Percival and Barnes, 2008) and pear (Raese, 1997). All the cold acclimated samples in our study (including plus- and minus-calcium) had higher freezing stress resistance than non-acclimated onions (Figure 4.2). There was no significant difference in freezing stress resistance between ACC and ACC+Ca with  $\text{LT}_{50}$  values around  $-25^\circ\text{C}$ ,  $-26^\circ\text{C}$ , respectively. However at lower temperature ( $-30^\circ\text{C}$ ), the freezing stress survival of ACC+Ca was 44% higher than ACC alone (Figure 4.2). The positive role of calcium during cold acclimation are also shown by studies where the applications of EGTA ( $\text{Ca}^{2+}$  chelator), and  $\text{La}^{3+}$  and verapamil ( $\text{Ca}^{2+}$  channel blockers) inhibit protein phosphorylation, development of freezing tolerance, and reducing the accumulation of specific mRNAs in alfalfa during cold acclimation (Monroy *et al.*, 1993). In red onions (*Allium cepa* L. cv Big Red), calcium is known to moderate the injury to the plasma membrane from freezing stress, and the application of exogenous calcium can also prevent cell plasmolysis and protoplasmic swelling resulting from the high concentration of extracellular  $\text{K}^+$  in freezing stress (Palta and Li, 1980; Arora and Palta,

1988; Palta, 1996). Mechanical stress, including mechanical deformation of the plasma membrane and the phase transition of the lipid compositions from the liquid to the solid state in plasma membrane (Yamazaki *et al.*, 2009) is considered to be the main cause of cell injury, which can be induced by the dehydration from freezing, rehydration from thawing, and ice crystal growth during the whole freeze-thaw cycles (Levitt, 1980). In research by Yamazaki *et al.* (2008), the freezing tolerance to mechanical stress in *Arabidopsis thaliana* is enhanced after cold acclimation by the extracellular calcium at both the cellular and tissue levels. Calcium is likely to contribute to the structural integrity of cell walls through the formation of cross-links between carboxyl groups of adjacent polyuronide chains (Steponkus, 1984). According to our data in Figure 4.2, exogenous application of calcium is capable of improving the freezing resistance regardless of whether the cells are non-acclimated or cold acclimated. Low temperature acclimation also results in an increase of calcium levels in the cytoplasm of plants, which improves the ability of plants to better respond in the temperature-sensing mechanism which then initiates the signal transduction pathway of cold acclimation (Minorsky and Spanswick, 1989; Knight *et al.*, 1996). At low temperature, the cold stimuli can trigger a series of changes including protein phosphorylation, cold acclimation-specific gene expression and freezing tolerance development (Monroy *et al.*, 1993; Chinnusamy *et al.*, 2007; Preston and Sandve, 2013). Collectively, these results suggest that calcium is involved in the acquisition of freezing resistance.



**Figure 4.2** Effect of cold and calcium on freezing hardiness. The survival rate of non-acclimated cells (NA), non-acclimated cells plus  $\text{CaCl}_2$  (NA+Ca), cold acclimated cells (ACC), cold acclimated cells plus calcium (ACC+Ca) plants was shown above. Error bars are means  $\pm$  SD based on five separate experiments.

Although we have demonstrated increased freezing resistance of Japanese bunching onion after cold acclimation and calcium treatment, it is not known how the treated cells resist freezing damage. Our hypothesis is that the cell wall is serving as a barrier to intracellular freezing. In order to test the hypothesis, it is necessary to understand the freezing process.

#### **4.2 Analysis of Freezing Process**

Using a cryostage connected to a Nikon ECLIPSE E400 microscope and 40X objective with video recording, the temperature at which ice nucleation occurred in the extra- and intracellular space of tissues was monitored during freezing. Subsequently, cell viability after thawing to 24°C was also observed (Table 4.1, 36 direct observations, see Appendix D for the complete set of data).

To study the ice-nucleation process, non-acclimated (plus- and minus-calcium) and two-week cold acclimated epidermal layers were incubated in the presence and absence of an extracellular nucleating agent, AgI. The application of exogenous ice nucleators plays an effective and vital role in triggering ice generation at a relative high temperature (-2 to -5°C) (Table 4.1). In our system, there was no spontaneous ice nucleation of the intracellular space in any of the > 40 samples tested across all treatments (Table 4.1). Thus, intracellular ice nucleators either do not exist or are inactive in the epidermal layer of onion. When intracellular nucleation did occur, this process was due to ice propagation from the extracellular space (apoplast) into the intracellular space (see the video in Appendix C). In the absence of artificial extracellular nucleation, freezing of non-acclimated cells (NA) spontaneously initiated in the extracellular space around -8°C and was immediately followed by intracellular ice nucleation with ice penetration through the apoplast into the intracellular space (Table 4.1). In the NA cells, there was no significant difference between the nucleation temperature of the extracellular and intracellular space. However, when artificial extracellular nucleation was induced around -4°C in NA+Ca cells, the temperature at which subsequent intracellular ice nucleation occurred (caused by propagation of ice across the apoplast into the cell) was delayed to around -8°C (Table 4.1). Without artificial extracellular nucleation, calcium treatment added to NA cells did not delay ice propagation from the extracellular to the intracellular space. However, the combination of calcium treatment and artificial extracellular nucleation (NA + Ca) (+) almost completely prevented ice propagation across the apoplast, and prevented intracellular ice nucleation. In this

case, the intracellular space of NA samples supercooled and the cells were alive to at least -10°C as determined by re-warming the cryostage to +24°C, and observing in protoplast streaming.

**Table 4.1** Summary of extracellular and intracellular freezing temperature in NA, ACC and NA+Ca treated plants

Treatment	AgI extracellular nucleating agent	Mean extracellular nucleation temperature (°C)	Mean intracellular nucleation temperature (°C)
NA	-	-8.2±1.1	-8.3±1.1
NA	+	-4.1±2.2	-8.4±1.2
ACC	-	-10.1±2.3	-15.1±4.8
ACC	+	-3.2±0.9	< -21.8±2.4
NA+Ca	-	-9.6±1.8	-9.6±1.7
NA+Ca	+	-3.1±0.8	< -20.7±3.3

NA: non-acclimated cells, ACC: cold acclimated cells,

NA+Ca: non-acclimated cells plus CaCl<sub>2</sub>

- no extracellular AgI nucleating agent added


+ extracellular AgI nucleating agent added

Data represents a total of 36 separate experimental observations.

The initial nucleation and growth of ice crystals in the extracellular space drives cellular water to the extracellular space from the protoplasm due to the low external cell water potential caused by the extracellular freezing, resulting in cell dehydration (Toivio-Kinnucan and Stushnoff, 1981). Once ice propagates across the cell wall and plasma membrane, intracellular freezing occurs, which is fatal to all biological cells (Fujikawa *et al.*, 1999). In nature, plant tissues can avoid intracellular freezing by tolerating freeze desiccation (Burke *et al.*, 1976). In our study, the combination of calcium with artificial extracellular ice nucleation changed the NA freezing process into extracellular freeze dehydration stress similar to the freezing process of highly acclimated samples. Generally, the freezing processes involve seven types (Gusta *et al.*, 2009) which were summarized in the Literature Review part of this thesis. One of the mechanisms involves supercooling in which the cells are not frozen below the freezing point. Survival of supercooled tissues at the temperature of liquid nitrogen (-196°C) has been observed in some temperate perennial plants (Burke *et al.*, 1976; Lipp *et al.*, 1994).

Without artificial nucleation, the freezing temperature of the intracellular space was only decreased by about 5°C (-10°C extracellular freezing temperature to -15°C intracellular freezing temperature) and the intracellular freezing temperature was not significantly different than the extracellular freezing temperature (Table 4.1). Similar to the NA + Ca and artificial extracellular nucleation, the ACC treatment in combination with artificial extracellular nucleation (at -3 to -4°C) resulted in almost complete prevention of ice propagation across the cell wall and plasma membrane. Virtually no intracellular ice nucleation was observed. Therefore, a barrier to avoid the growth of ice crystals must exist in all types of freezing processes under either avoidance or tolerance strategies. This barrier is able to not only prevent ice propagation into a cell or from stems into flowers and fruits but also decrease the loss of intracellular water because of extracellular freezing (George and Burke, 1977; Ashworth and Abeles, 1984; Wisniewski and Davis, 1995; Workmaster *et al.*, 1999). Ice crystals in rhododendron and *Holcus lanatus* leaves are formed first on the leaf surface and the growth of ice is delayed through this area (Wisniewski and Fuller, 1999; Pearce and Fuller, 2001). Ice is formed and propagated from the cortex to the xylem in peach stem (Wisniewski *et al.*, 1997). The thick cuticles and xylem ray parenchyma cells play critical roles in preventing ice propagation into the cells as the barrier.

The epi-cuticular layer (Figure 4.3) is the first physical barrier to prevent ice propagation into the cells of the whole plant. The epi-cuticular layer is attached to the epidermal cells through the middle lamella and cell wall. The properties such as porosity and permeability of the cell wall are thus essential factors affecting extracellular ice formation. Once ice crystals are generated in the extracellular space, the risk of cell dehydration increases and finally cell collapse occurs (Rajashekar and Burke, 1996). With cold acclimation, the thickness of leaf cell walls increases, protecting the cells against intracellular freezing (Griffith and Brown, 1982; Stefanowska *et al.*, 1999). Under abscisic acid-induced freezing tolerance, cell wall thickness is also significantly increased and the majority of externally added <sup>14</sup>C-sucrose incorporates into this insoluble fraction over time, paralleling the enhancement of freezing tolerance (Tanino *et al.*, 1991). In our study, we provide direct visual evidence that intracellular freezing was delayed or did not occur in the cold acclimated onions, which may be related to compositional and structural changes of the onion cell walls. To our knowledge, this is the first time that such visual observation of ice propagation from the extracellular space to nucleate the intracellular space has been recorded.



**Figure 4.3** Organization of the plant cuticle (Kourouniotti *et al.*, 2013). Epidermal cells produce the cuticle, a polymer matrix known to contain cutin, waxes and sometimes polysaccharides (pectin) of the cell wall.

As mentioned, there was no spontaneous intracellular freezing prior to the extracellular freezing in any of these NA and ACC samples, no matter how low the temperature of supercooling. Thus, there do not appear to be active ice nucleators which initiate intracellular freezing at a high (less negative) temperature. Without artificial nucleation, extracellular freezing initiated at relatively low temperatures (average of  $-9^{\circ}\text{C}$ ) in all treatments and thereafter tissues supercooled to relatively low temperatures before freezing. In instances when supercooling occurred (no freezing) and tissues were re-warmed from this supercooled state, cells were observed to be viable through active protoplasmic streaming (see Appendix D). However, once extracellular freezing spontaneously initiated at low temperatures in the supercooled state, intracellular freezing immediately followed in all cases (19 out of 19 samples, NA, NA+Ca and ACC) with the average subsequent intracellular nucleation temperature of  $-11^{\circ}\text{C}$ . Not surprisingly, all these cells were dead upon thawing. Acclimated tissue did not induce higher temperatures of extracellular nucleation compared to NA and thus acclimation itself did not prevent supercooling of the extracellular space. Also as expected, in the presence of the artificial nucleator in the extracellular apoplastic space, freezing was induced in this region at relatively high temperatures (average of  $-4^{\circ}\text{C}$ ). Under these artificial extracellular nucleation conditions, subsequent intracellular freezing was either delayed (in NA samples, average of  $-8^{\circ}\text{C}$ ) or did not

occur (NA+Ca and ACC). The combination of AgI nucleation and calcium treatment had a pronounced effect on virtually eliminating intracellular freezing even in NA samples. Olien (1961) indicated that the lower the temperature of freezing, the higher the physical force (biophysical force of ice propagating throughout the tissue), which can easily cause injury to the tissue.

Calcium is a divalent cation ( $\text{Ca}^{2+}$ ) and plays various roles in plant growth and development, such as a macronutrient element in the whole plant, an intracellular secondary messenger in the cytosol, and as a structural integrity support in the membrane and cell wall (Wyn Jones and Lunt, 1967; Emanuelsson, 1984; Palta, 1990; Schiefelbein *et al.*, 1992; Palta, 1996; White, 1998; White and Broadley, 2003; Hepler, 2005). Calcium exists in the cell wall by forming  $\text{Ca}^{2+}$  pectate with acidic pectin residues which are de-esterified by pectin methylesterase (PME) (Hepler, 2005; Tuteja and Mahajan, 2007). More specifically, during the formation of the cell wall, pectin releases the carboxyl groups which bind  $\text{Ca}^{2+}$  under the catalysis by PME, thus generating gel-like cross-linkages between cells. This process increases cell wall gelation and forms the numerous small pores in the cell wall. The increased gelation has the ability to maintain the integrity of the cell wall, which provides protection against freezing damage since the cell wall can be the primary physical barrier. Moreover, the combination of AgI extracellular nucleation and ACC treatment completely eliminated subsequent freezing of the intracellular space. Ice movement from the extracellular space to the intracellular space appeared to be blocked by the apoplast under these treatment combinations. Cells remained viable after extracellular nucleation to at least  $-10^{\circ}\text{C}$  before intracellular nucleation occurred (Appendix D). However, even in the absence of intracellular freezing, with prolonged extracellular freezing, both NA+Ca and ACC cells were killed by extracellular freeze dehydration stress when observed at the limit of the cryostage ( $-23^{\circ}\text{C}$ ). Thus, the blockage of ice alone cannot prevent injury in freezing tolerant cells. It can, however, play a significant role in relatively unhardy tissue and in those organs which survive through freezing avoidance mechanisms. Nevertheless, greater gelation of the apoplast will slow water loss during extracellular freeze dehydration stress and will also slow rehydration during thawing. These two factors may play a role in increasing freezing stress resistance in tolerant cells and should be investigated.


Although the freezing process (extra- and intra-cellular freezing) has been well documented, the process by which ice crystals pass through the cell wall is less studied. Thus,



the specific biochemical changes and regulation of these changes occurring in the apoplast to prevent ice penetration to the protoplasm after cold acclimation should be examined.


### **4.3 Studies of Apoplastic Permeability**

Differential cell wall permeability was observed in non-acclimated and acclimated samples using a combination of fluorescent dyes and confocal microscopy. Fluorescence was observed in both the apoplast and cytoplasm of non-acclimated samples which indicated that the dextran conjugated to fluorescein penetrated into the intracellular space of non-acclimated samples up to the tonoplast (Figure 4.4a). This phenomenon was less pronounced in the acclimated sample (Figure 4.4b). Figures 4.4c and d are visible light images of the cells in Figures 4.4a and b, respectively. The pore diameters of a 70,000 and 3,000 MW dextran are about 12 and 2.6 nm, respectively (Wang and Fisher, 1994; Zhang *et al.*, 2011). The smaller the pores of the cell wall, the less ice crystal penetration through the cell wall. In our case, the pore size of acclimated cells was reduced down to at least 2.6 nm since the 3,000 MW dextran could not penetrate the acclimated cell walls. In grape cells, the cell wall pore size was decreased from 3.5 nm to 2.2 nm. Similarly, cold acclimation in apple reduces the pore size from 2.9 nm to 2.2 nm after cold acclimation (Rajashekar and Lafta, 1996). Jones *et al.* (2000) measured the apoplastic permeability with fluorescent rhodamine green dyes and found the dyes in non-acclimated *Vitis vinifera* cv. Chardonnay buds could penetrate through the primordia because of the permeable bud axis. However, it did not occur in the acclimated buds since a permeability barrier developed that prevented dye propagation. This is consistent with the decreased permeability of the acclimated onion cells, which supports the role of the cell wall in the epidermal onion layer as a barrier.



**Figure 4.4** The impact of acclimation on apoplast permeability. The penetration of fluorescein conjugated with dextran (3,000 and 70,000 MW) between non-acclimated (NA) and acclimated (ACC) was measured using a confocal microscope (50X). a: non-acclimated sample; b: 2 week acclimated sample; c: non-acclimated sample with white light; d: 2 week acclimated sample with white light. Scale bar = 50  $\mu\text{m}$ .

In order to evaluate how small the pore size might be, we used the smallest known diameter fluorescent molecule available, fluorescein without conjugation which has an effective radius of 6.5 Å (equal to 1.3 nm in diameter) (Lawrence *et al.*, 1994). The diffusion rates of fluorescein across the cell wall to the intracellular space between NA and ACC were measured based on the brightness of fluorescence with a confocal laser scanning microscope (LSM) and NIS-Elements software. According to the video from the LSM (Video in the Appendix E), the fluorescein penetrated into the NA cells almost immediately after application to the extracellular space. By contrast, the fluorescein in the ACC samples was either markedly delayed or did not penetrate at all the cells. In Figure 4.5, the calculated mean diffusion rate in NA was 28.95 pixels per second, which was 70 times faster than in ACC cells (0.41 pixels per second).



**Figure 4.5** The diffusion rate of fluorescein in non-acclimated (NA) and cold acclimated (ACC) cells. Error bars are means of 5 replications  $\pm$ SD.


Fluorescein is a pH dependent dye and the fluorescence can be influenced by different pH conditions (Zhu *et al.*, 2005; Lakowicz, 2006). The fluorescent density of fluorescein increases when the pH value increases from 6.9- 8.4, but it almost remain unchanged when pH value is greater than 8.4 (Zhu *et al.*, 2005). Therefore, the cellular pH may also affect the measurement of diffusion rate. After cold acclimation, the onion cells may be more acidic because some pectins become hydrolysed to acidic pectin and methanol by the catalysis of PMEs. Thus, the observed fluorescence in acclimated cells should be reduced compared to non-acclimated cells. In other words, the real diffusion rate in acclimated cells may be greater than the calculated value above. According to the study from Carpita *et al.* (1979), the cell wall pores in living plant cells are approximately 5 nm in diameter. In un-extracted onion cell wall material, the diameter of most pores is less than 10 nm and the effective pore size *in vivo* becomes smaller because of the existence of bound water (McCann *et al.*, 1990). In our cold acclimated onion cells in the majority of samples, fluorescein did not penetrate into the acclimated cells. The pore size of the onion cell walls after cold acclimation decreased, to less than 1.3 nm diameter based on the diameter of pure fluorescein. The diameter of water is around 0.27 nm at 25 or 40°C (Schatzberg, 1967), while the diameter of supercooled water is only about 9  $\mu$ m at -46°C (Sellberg *et al.*, 2014). In non-acclimated onion cells, water can freely flow in and out of the cells. Conversely,

this phenomenon may be restricted in the cold acclimated cells as a consequence of smaller pore size of the cell wall. Ashworth and Abeles (1984) also indicated water freezes at very low temperatures when contained within small pores (less than 100 nm). According to the melting point curve, water was assumed to freeze at -15 to -25 °C when the pore size of capillaries is 4 nm in diameter (Ashworth and Abeles, 1984). The diameter of pores in non-acclimated onion cell walls is between 0-10 nm, so the freezing temperature was predicted to be slightly higher than -15°C. In contrast, the freezing temperature in cold acclimated onion cells is expected to be much less than -25°C since the pore size in our acclimated onion cells is less than 1.3 nm diameter (according to fluorescein tests). These smaller wall pores may effectively reduce the freezing point of intracellular water and prevent the growth of ice crystals into the cells. The cell wall permeability is affected by the cell wall porosity which is regulated by the pectin gelation. Pectin gelation can be modulated by the degree of pectin methyl-esterification and the activity of pectin methylesterases (PMEs). Therefore, both the degree of pectin methyl-esterification and the activity of pectin methylesterases (PMEs) in Japanese bunching onions were measured in the following experiments.

#### **4.4 Analysis of The Degree of Pectin Methyl-esterification**

To study the degree of pectin methyl esterification, the monoclonal antibodies JIM 5 and JIM 7 were used (Figure 4.6). Monoclonal antibody JIM5 binds to partially methyl-esterified epitopes of homogalacturonic acid (HG) and can also bind to un-esterified HG; and JIM7 binds to partially methyl-esterified epitopes of HG but does not bind to un-esterified HG (Knox *et al.*, 1990; Clausen *et al.*, 2003; Verhertbruggen *et al.*, 2009). The fluorescence of JIM5 HG epitopes in ACC ligule cells (Figure 4.6b) was distinctly stronger than in NA control cells (Figure 4.6a). With JIM7 application, there was a slight reduction in fluorescence intensity in the ACC treatments compared to NA (Figure 4.6c and d). These data were initially obtained from Norwegian University of Life Sciences Imaging Lab, Ås, Norway and later confirmed at the University of Saskatchewan where the same experiment was repeated in NA and ACC Japanese bunching onion epidermal layers with cross sections. A stronger signal of JIM5 and weaker signal of JIM7 in acclimated cells from epidermal layer was noted when compared to the non-acclimated cells (see APPENDIX G). The stronger signal of JIM5 in the cold acclimated onions indicates the un-esterified pectins increased after cold acclimation, which may help to form more gelation through calcium cross-linkage, thereby decreasing the cell wall permeability and

preventing the growth of extracellular ice crystals into the cells. By contrast, the weaker signal of JIM7 showed that the esterified pectins decreased after cold acclimation, which may have hydrolysed to un-esterified pectin and methanol. These results were also supported by other studies in which the labeling with JIM 5 is more noticeable than JIM 7 fluorescence in December peaches and cold-acclimated winter oil-seed rape plants, respectively (Wisniewski and Davis, 1995; Solecka *et al.*, 2008).



**Figure 4.6** Immunohistochemical studies of pectin composition in non-acclimated (NA) and acclimated (ACC) epidermal layer cells with monoclonal antibodies JIM 5 and JIM 7. a: non-acclimated cells with JIM 5, b: cold acclimated cells with JIM 5, c: non-acclimated cells with JIM7, d: cold acclimated cells with JIM7. Scale bar = 100 µm.

The apoplastic space is composed of the region outside of the plasma membrane which includes the cell wall and middle lamella. It plays significant roles in the interaction between the plant and the environment. The cell wall provides a primary natural barrier to prevent damage from the hostile environment to the plant. As one of the major cell wall components in Type I cell walls, pectin forms a hydrated cross-linked three-dimensional network in the matrix of primary plant cell walls and interlocks with hemicellulose to help to bind adjacent cells together (Carpita and Gibeaut, 1993; Voragen *et al.*, 2009). It also regulates the porosity and permeability of the cell wall (Baron-Epel *et al.*, 1988; Wolf *et al.*, 2009; Peaucelle *et al.*, 2012) and provides a defense against pathogens such as powdery mildew, in which pectin accumulation is released by


the hydrolytic enzymes (pectate lyase), eliciting the defense response to mildew (Vogel *et al.*, 2002; Vorwerk *et al.*, 2004). Pectins also appear to be involved in drought stress in which cell walls in the albedo of immature lemon fruitlets are thickened and its high water-binding capacity is maintained by the high content of pectin (Schröder *et al.*, 2004). Under freezing stress, resistance was improved by modifications in the degree of pectin methyl esterification (Wisniewski and Davis, 1995; Solecka *et al.*, 2008). In Wisniewski and Davis's study (1995), the labelling with JIM7 recognizing highly esterified epitopes of pectin was distributed in the whole pit membrane, primary cell walls of xylem and floral bud tissues of peach, while the labelling with JIM5 recognizing un-esterified epitopes of pectin was located outside of the pit membrane in xylem tissues and in middle lamellae of floral buds. These results revealed the distribution and types of pectin in both xylem and floral bud tissues of peach. It has been known that deep supercooling occurs in the xylem tissues and floral buds in peach. Combining the results above, pectin in the xylem tissues and floral buds greatly contributes to deep supercooling, avoiding freezing damage. Meanwhile, JIM5 signal in December peaches was stronger than in July/August samples, indicating there were more un-methylesterified pectins in December peaches which could survive in the winter. Pectins are involved in enabling freezing avoidance to occur in xylem ray parenchyma of deep supercooling trees through ice blockage by the xylem pits (Ashworth and Abeles, 1984; Wisniewski *et al.*, 1991a). However, the study from Baldwin *et al.* (2014) indicated higher immunolabeling with JIM7 in the frost-tolerant pea genotype 'Champagne' than the frost-sensitive pea genotype 'Terese', which indicates an increase in the degree of methyl-esterification of pectins after cold acclimation.

## **4.5 Cell Wall Compositional Analysis**

### **4.5.1 Determination of Cell Wall Quantity in Ligule Tissue**

The relative crude cell wall preparations of the epidermal layer, non-epidermal layer and the whole sheath (composed of both epidermal layer and non-epidermal layers) during cold acclimation were analyzed. The epidermal layer contained significantly higher relative cell wall content (insoluble fractions), occupying up to 22.6 % of fresh sample weight after 14 days cold acclimation compared with NA onion epidermal peels (Figure 4.7). The relative cell wall content in the non-epidermal layers and the whole sheath increased slightly during cold acclimation. It has been shown that the thickness of the cell wall increases significantly in the hardened winter

rye (Griffith *et al.*, 1985) and in the 7-days ABA-acclimated bromegrass cells (Tanino *et al.*, 1991). The cell wall content in winter oil-seed rape leaves also increases after cold hardiness (Kubacka-Zębalska and Kacperska 1999; Solecka *et al.* 2008), which is consistent with the result from the ACC onion cell walls. The research from Stefanowska *et al.* (1999) illustrated increased cell wall content was related to the increased thickness of cell wall in epidermis cells. The changes of the cell wall structure were dependent on cold hardiness and may have contributed to the protection from freezing-induced damage since these modifications increase leaf tensile stiffness which avoids cell volume collapse and cell wall deformation in response to extracellular freezing (Pearce, 1988; Fujikawa *et al.*, 1999; Stefanowska *et al.*, 1999; Jarvis and McCann, 2000).




**Figure 4.7** The relative content of onion crude cell wall mass during cold acclimation. Values are expressed by the percentage ratio of the crude cell wall fresh weight of respective tissue. e: epidermal layer, n: non-epidermal layer, w: whole sheath (e+n). Error bars are  $\pm$  SD of seven separate experiments.

#### 4.5.2 Cell Wall Carbohydrate Analysis of Ligules

The individual cell wall components in the epidermal layer were quantified by HPLC (GC/MS) analysis and partitioned into TFA insoluble (trifluoroacetic acid) and water soluble fractions. As expected, the insoluble TFA fraction contained a higher concentration of cell wall components than the water soluble fraction. The cell wall polysaccharides of onion contain fucose, rhamnose, arabinose, galactose, glucose, xylose, mannose, galacturonic acid and glucuronic acid (Ng *et al.*, 1998). The major components within the onion epidermal layers were

galactose and galacturonic acid, representing 28.5% and 35.5% of the total cell wall fraction, respectively (Figure 4.8). Galacturonic acid (GalA) including methylated and un-methylated residues is the main component of pectin, representing around 70% of homogalacturonan (Mohnen, 2008). Carbohydrate composition was similar between NA and ACC except for galacturonic acid in which the ACC samples contained slightly higher about 27% galacturonic acids than non-acclimated onion cells in the insoluble fraction. Pectic polysaccharides are the major group of onion cell wall constituents, occupying 42.2% of the cell wall (Mankarios *et al.*, 1980). The higher content of GalA in the cold acclimated onion cells implies there was more pectin in acclimated onion cells than in the non-acclimated onions. This is consistent with the studies of pea (Weiser *et al.*, 1990), winter oil-seed rape plants (Kubacka-Zębalska and Kacperska, 1999; Stefanowska *et al.*, 1999; Solecka *et al.*, 2008) and the tolerant pea cultivar ‘Champagne’ (Baldwin *et al.*, 2014) during the period of cold acclimation.



**Figure 4.8** Cell wall components of the epidermal layer. Data were obtained by analysis of GC/MS NA and ACC of epidermal ligule cells. The cell wall components include trifluoroacetic acid (TFA) fraction (insoluble) and water soluble fraction. NA: non-acclimated onion cells, ACC: cold acclimated cells. Error bars are  $\pm$  SD of two separate experiments. Fuc: fucose; Rha: rhamnose; Ara: arabinose; Gal: galactose; Glc: glucose; Xyl: xylose; Man: mannose; GalA: galacturonic acid; GlcA: gluconic acid.


## 4.6 PME Activity in Epidermal Tissues of Ligules

### 4.6.1 Determination of Total Cell Wall Protein Content

The relative quantity (% cell wall protein/total crude cell wall) of total cell wall protein in the epidermal layer was two to three times greater in the epidermal layer than in the non-epidermal layer in both NA and ACC tissue (Figure 4.9). Specifically, the total cell wall protein occupied 4% of the total crude cell wall after 14 days cold acclimation, which was significantly




greater than the non-epidermal layer and the whole sheath (Figure 4.9). However, the rate of cell wall protein increase appeared to be similar between tissues over 14 days of acclimation (Figure 4.9). The plant cell wall protein profile includes in decreasing order of concentration: antifreeze proteins (Duman and Olsen, 1993; Griffith *et al.*, 2005; Yu *et al.*, 2010), arabinogalactan proteins (AGPs) (Gong *et al.*, 2011), expansins (Noh *et al.*, 2009), glycine-rich proteins (GRPs) (Ferullo *et al.*, 1997), xyloglucan endotransglucosylase/hydrolases (XTHs) (Yun *et al.*, 2007), extensin (Weiser *et al.*, 1990), and related cell wall enzymes like pectin methylesterases (Kubacka-Zębalska and Kacperska, 1999; Solecka *et al.*, 2008; Baldwin *et al.*, 2014). This profile may change during cold acclimation and these proteins may up- (AGP, expansins) or down-regulate (extension and XTHs) gene expression, modifying the mechanical properties of the cell wall such as mediating cell-wall loosening by expansins (Li *et al.* 2003) and determining the cell wall plasticity by XTH (Chanliaud *et al.*, 2002), thereby potentially improving the freezing tolerance of plants (Yun *et al.*, 2007; Noh *et al.*, 2009; Shi *et al.*, 2014; Gall *et al.*, 2015).



**Figure 4.9** Relative total cell wall protein content during cold acclimation. Values are expressed by the percentage ratio of the total cell wall protein content to the crude cell wall FW. e: epidermal layer, n: non-epidermal layer part, w: whole sheath. Error bars are  $\pm$  SD of seven separate experiments.

#### 4.6.2 PME Activity Analyses

Pectin methylesterases (PMEs, EC 3.1.1.11), a cell wall-associated enzyme in the apoplast, catalyses the hydrolysis of the methyl ester group of pectin (mainly homogalacturonan) to release acidic pectin and methanol; the carboxyl acid residues from this reaction can be cross-linked by calcium, forming calcium-pectate gels, which can increase the stiffness of the cell wall (Jarvis, 1984; Bosch, 2005). PME activity (expressed as  $\text{nEq H}^+ \text{mg}^{-1} \text{protein min}^{-1}$ ) in the cell wall was separated into the soluble and cell wall-bound enzyme fractions (Figure 4.10). Overall, the PME specific activity in the soluble fraction followed a decreasing tendency over the 14 day ACC period (Figure 4.10a). By contrast, PME had an increasing tendency in the cell wall-bound fraction following the acclimation period (Figure 4.10b). The relative PME enzyme activity in the epidermal layer was significantly higher than the non-epidermal layer and the whole sheath. Specifically, after two weeks of cold acclimation, the relative PME activity in the epidermal layer (expressed as a function of each tissue's respective fresh weight) was more than two times that of the non-epidermal layer (Figure 4.10c).




**Figure 4.10** Relative PME specific activity during cold acclimation. The PME activity unit was expressed by  $\text{nEq H}^+ \text{mg}^{-1} \text{protein min}^{-1}$ . a: soluble PME activity, b: wall-bound PME activity, c: total PME activity (including soluble and wall-bound PME activity), e: epidermal layer, n: non-epidermal layer part, w: whole sheath (e+n). Error bars are means of five replications  $\pm$  SD.

There are two main types of primary cell wall among species of angiosperms: Type I cell walls from eudicots and some monocots, and Type II cell walls from the Poaceae. In Type I cell walls, the network of cellulose microfibrils is tethered with xyloglucan polymers and embedded in a gel of pectic polysaccharides through the cross-linking with calcium. By contrast in the Type II cell walls, the microfibrils are interlaced with glucuronoarabinoxylans rather than xyloglucans, and there is rarely pectin (Carpita and Gibeaut, 1993). The cell walls of onions belong to the Type I cell wall which contain abundant pectic matrix (Carpita and Gibeaut, 1993). Pectins impart various physical characters to the cell walls, such as density, wall porosity, the distribution of wall enzymes and proteins, and cell-to-cell adhesions (Albert *et al.*, 2002). The methylester groups of cell wall pectins are hydrolysed to acidic pectin and methanol through de-methylesterification by PME (O'Neil *et al.*, 1990; Pelloux *et al.*, 2007). The acidic pectins with free carboxyl groups are capable of forming cross-links with calcium, increasing the gelation of pectin, thereby reducing the pore size and permeability of cell walls. The ability of cell walls to resist extracellular ice nucleation into the cells may be improved through the modification of the cell wall structure. There is a negative correlation between homogalacturonan (HG) methyl esterification and PME activity in chilling-injured peach fruit after storage at 5°C for 1 to 4 weeks, in that increased PME activity reduces the accumulation of methyl esterified HG, which increases cold storage of peach (Brummell *et al.*, 2004). Un-methylesterified HG can form a gel through cross-linkage with calcium, which in turn may regulate the freezing resistance by enhancing the formation of gelation and cell wall porosity (Willats *et al.*, 2001), thereby potentially decreasing the risk of ice penetration into the cells. PME activity in oil seed rape plants increases after cold acclimation (Solecka *et al.*, 2008), which is consistent with our result in cold acclimated onions. Cell wall rigidity is also affected by de-methylesterification of HG (Willats *et al.*, 2001; Solecka *et al.*, 2008). The plant cell walls play a critical role in withstanding the hydraulic turgor pressure of the cells, which combine with wall construction to contribute to cell wall rigidity (Lodish *et al.*, 2000). During freezing stress, extracellular ice nucleation induces intracellular dehydration stress which lowers internal water potential and leads to the decrease in the turgor pressure. Increased cell wall rigidity accompanied by solute accumulation helps to maintain cell volume at low water potentials during drought stress (Clifford *et al.*, 1998). Thus, cell wall rigidity increases in response to lowering water potential under intracellular dehydration stress. Ha *et al.* (1997) measured the cell wall rigidity in dry and hydrated onions with solid-state

nuclear magnetic resonance relaxation and found that both pectic matrix and microfibrils provide mechanical properties and rigidity to the cell walls. In addition, the transfer from the esterified pectins in the apical pollen tube to un-esterified pectins in the distal pollen tube was thought to be associated with the increase in the cell wall rigidity (Parre and Geitmann, 2005). The overall cell wall rigidity appears to result from the un-methylated pectins (Fenwick *et al.*, 1997). After cold acclimation, the amount of un-methylated pectins increased (Fig.4.6) mainly due to the higher PME activity comparing the non-acclimated cell walls. More un-methylated pectins mean more rigid cell walls. Therefore, the rigidity of the cell wall is improved by the increasing PME activity, and this is accompanied by the ability of cell wall to resist freezing damage.

As compared with common onions (*A. cepa*) and green onions (*A. chinense*), Japanese bunching onion (*A. fistulosum*) had higher PME activity in soluble, bound and total fractions (Figure 4.11). *A. cepa* is very sensitive to low temperature and has a low freezing resistance with -11°C of LT<sub>50</sub> value (Palta *et al.*, 1977), while LT<sub>50</sub> in *A. fistulosum* is -27°C. The high PME activity in *A. fistulosum* may explain why Japanese bunching onion has the highest freezing resistance capability among these three species.



**Figure 4.11** The PME activity in the whole sheath (including epidermal layer and non-epidermal layer) of onion species: Common onion (*A. cepa*), green onion (*A. chinense*) and Japanese bunching onion (*A. fistulosum*) were analyzed. Error bars are means of four separate experiments  $\pm$  SD.

PME also plays a vital role in dormancy breakage (Ren and Kermode, 2000), seed germination (Ren and Kermode, 2000; Müller *et al.*, 2013), elongation growth (Bordenave and Goldberg, 1993; Peaucelle *et al.*, 2012), plant development (Wolf *et al.*, 2009), fruit ripening

(Kagan-Zur *et al.*, 1995; Draye and van Cutsem, 2008), pathogen defense (Lionetti *et al.*, 2007; Volpi *et al.*, 2011). PME activity affects the plant-pathogens interactions (Micheli, 2001; Lionetti *et al.*, 2012), but the way in which PME acts on plant pathogen resistance is opposite from freezing stress. As one of the cell wall degrading enzymes (CWDEs), PME affects the degree of pectin methyl esterification. A series of studies have revealed that the resistance to *Pectobacterium carotovorum* in potato cultivars (McMillan *et al.*, 1993; Marty *et al.*, 1997), *Colletotrichum lindemuthianum* in resistant bean cultivars (Boudart *et al.*, 1998) and *Ralstonia solanacearum* in tomato genotypes (Wydra and Berl, 2006) are correlated with higher degree of pectin methyl esterification. Therefore, in contrast to the function of PME activity in freezing stress, the reduced susceptibility of plants to the various pathogens is dependent on the low PME activity and high degree of pectin methyl esterification (Lionetti *et al.*, 2007; Lionetti *et al.*, 2012).

In our study, the Japanese bunching onion was used as a model system to better understand the mechanisms of intracellular freezing stress avoidance based on cell wall alterations. The LT<sub>50</sub> decreased from -12°C to -15°C in one-week acclimated samples and significantly lowered to -27°C after two weeks of cold acclimation. Freezing resistance of Japanese bunching onion cells increased greatly after two-week acclimation compares to one-week acclimation. Based on the measurements on the apoplast compositions during cold acclimation, the crude apoplast contents increased by 45% from non-acclimation to one-week acclimation while it slightly increased by 9% from one-week to two-week acclimation. However, the big increase of PME activity occurred from one-week (15%) to two-week (36%) acclimation, which was consistent with the increasing tendency in freezing resistance during the period of cold acclimation. This result indicated that the improving of resistance to freezing is more dependent on the efficiency of pectin methylesterase not only because of the increasing amount of the apoplast.

There is no doubt that cold acclimation plays a very important role in improving freezing resistance through modifications in morphology such as rosette shape in hardy varieties while upright shape in sensitive varieties of flax during winter (Omran *et al.*, 1968), changes in physiological and molecular level such as antifreeze proteins (AFPS) accumulation in the apoplast during cold acclimation (Griffith *et al.* 2005), changes in ultrastructure such as cell wall (Alonso *et al.*, 1997). In this study, we hypothesized that the alterations in cell wall composition

in response to low temperature regulates freezing avoidance through alterations in cell wall permeability. In the freezing process, ice nucleation first occurs in the extracellular space, and ice crystals may subsequently propagate through the cell walls and plasma membrane, nucleating the protoplast causing cell death. In tolerant cells, the collapse of the cell walls is due to extracellular freeze-induced cell dehydration which is related to water movement out to the extracellular space (Levitt, 1980; Rajashekar and Burke, 1996).

The cell wall porosity determines the permeability of cell wall to allow the molecules with certain molecular weight such as water and proteins to penetrate into the cells (Albersheim *et al.*, 2011). The lower porosity with smaller pore size of cell wall can restrict the penetration of the big molecules like ice crystals, reducing the freezing damage to the cells (Jones *et al.*, 2000). As the primary physical barrier, the mechanical properties of cell walls greatly contribute to the resistance of plants to freezing stress (Tao *et al.*, 1983; Murai and Yoshida, 1998) and are affected by the amount and structure of polysaccharides in the cell walls (Taiz, 1984; Carpita and Gibeaut, 1993). Generally, the primary cell walls in most plant cells are freely permeable and allow small molecules such as water (diameter of about 0.27 nm at 25°C) and some small proteins to pass through to the intracellular space. The mixture of 70,000 MW (12 nm in diameter) and 3,000 MW (2.6 nm in diameter) dextran conjugated with fluorescein in two-week cold acclimated (12/4°C day/night) Japanese bunching onion was blocked out of the cells while the dye penetrated into the tonoplast region in non-acclimated cells. This is consistent with the study from Jones *et al.* (2000) who illustrated that both the dye and ice crystals cannot pass through the grape bud axes due to the permeability barrier in the axis of buds after cold acclimation. Meanwhile, they also used pectinase to remove pectin, and found that the apoplastic permeability increased and ice nucleation temperature also increased. Therefore, pectins are involved in the permeability barrier to limit the ice propagation. According to our study, cold acclimation induced an increase in the amount of galacturonic acid (GalA), the basic component of pectins (mainly homogalacturonan). In addition, un-methylated pectin increased in the cold acclimated cells as observed using immunolabeling. Cross-linkage in the plant cell walls is formed through the ionic bond combination of anionic un-methylated pectin and cationic calcium, which promotes the production of pectate gel and provides cell wall porosity (Ridley *et al.*, 2001; Liu *et al.* 2009; Voragen *et al.*, 2009). In this case, the cell wall has the characteristic permeability to permit the passage of molecules having suitable diameter through cell wall pores.

The survival rate in the treatment of ACC+Ca was much higher than that in ACC alone, by approximately 44%. Calcium is involved in improving the freezing resistance in response to cold acclimation. Apoplastic  $\text{Ca}^{2+}$  is the highest concentration in plant tissues, about 60-75% of the total tissue  $\text{Ca}^{2+}$  content (Demarty *et al.*, 1984), playing an essential role in controlling cell integrity, cell wall cohesion and plasma membrane permeability (Hirschi, 2004). Studies have demonstrated that exogenous applications of calcium enhances tolerance to cold stress (Wang *et al.*, 2009; Zhou and Guo, 2009), which is consistent with my study showing increased survival rate after calcium application regardless of non-acclimated or acclimated cells. In addition, cytoplasmic calcium levels increased rapidly in both *Arabidopsis* (Knight *et al.*, 1996; Polisensky and Braam, 1996) and alfalfa (Monroy and Dhindsa, 1995) in response to low temperature, which is thought to be mainly due to an influx of calcium from extracellular stores. Cytosolic calcium has been shown to regulate the basic components of calcium signal transduction pathway, such as calmodulin (CaM), and calcium-dependent protein kinases (CDPKs), thereby improving the resistance to freezing stress since calcium serves as a second messenger in response to cold acclimation (Trewavas and Gilroy, 1991; Roberts and Harmon, 1992; Bush, 1993; Monroy *et al.*, 1993). Here, the roles of calcium in the cell wall during freezing stress were addressed. High content of apoplastic  $\text{Ca}^{2+}$  provides more opportunity to bind the contiguous un-methylesterified pectin (mainly homogalacturonan) to form the ‘egg-box’ structures which confers the rigidity of cell wall and maintains the integrity of cell wall in order to relieve the stresses (Grant *et al.*, 1973; Hamann *et al.*, 2009). In turn, the un-methylesterified pectins have a high affinity with calcium ions (Goldberg *et al.*, 1996). Moreover, the un-methylesterified pectin is released from the methylesterified pectin through the catalysis by pectin methylesterases (PMEs) which are a hydraulic enzyme residing in the cell wall (Micheli, 2001; Pelloux *et al.*, 2007). PMEs help to form very tight intermolecular interactions (pectate gel) with  $\text{Ca}^{2+}$  (Bosch, 2005), and regulate apoplastic  $\text{Ca}^{2+}$  content in response to stresses in order to modulate the pectic network (Micheli, 2001; Pelloux *et al.*, 2007). Therefore, PMEs and apoplastic  $\text{Ca}^{2+}$  interact with each other. Guillemain *et al.* (2008) showed PME plus  $\text{Ca}^{2+}$  increased the mechanical rigidity of the cell wall structures through a vacuum-impregnation experiment. After cold acclimation, PMEs activity increased in Japanese bunching onions, resulting in more un-methylesterified pectins, tighter gel formation and smaller pore size. The pore size of the cell wall determines its permeability (Baron-Epel *et al.*, 1988). In my study, I did



find cell wall permeability was reduced, according to the blockage of 3 000 MW dextran conjugated with fluorescein (2.6 nm diameter) and slower diffusion rate of pure fluorescein (1.3 nm diameter) in cold acclimated cells, thus the diameter of pore size of acclimated cell walls should be less than 1.3 nm. However, the diameter in non-acclimated cell wall pores is larger than 2.6 nm since the mixed dyes of 3,000 and 70,000 MW (12 nm in diameter) dextran conjugated with fluorescein can freely pass through non-acclimated cell walls. Furthermore, the immunolabeling test showed the stronger signal of JIM5 and the weaker signal of JIM7 appeared after cold acclimation, which further confirmed the un-methylesterified pectins increased in response to cold. Also, the increase in cell wall and pectin content may contribute to a thickened cell wall that provides a stronger barrier against ice penetration. The results from this project address my hypothesis well that alterations in cell wall compositions (pectin, especially un-methylesterified pectin) in response to low temperature improve freezing avoidance through a decrease in cell wall permeability which results from the tighter binding between un-methylesterified pectin and  $\text{Ca}^{2+}$ . Except freezing stress, it has been shown that some other stresses including drought and pathogen attacks also can be avoided through the modifications of cell wall (Micheli, 2001; Verslues *et al.*, 2006; de Freitas, *et al.*, 2012). Consequently, the cell wall is really a primary physical barrier to avoid environmental injuries.

## 5.0 CONCLUSIONS

It is well-known that intracellular freezing is lethal to all cells. As a freezing stress resistance mechanism, freezing avoidance provides plants with more potential to survive subzero temperatures without intracellular freezing. The hypothesis in this project that reduced permeability of the apoplast through pectin modification will increase freezing avoidance and freezing stress survival was supported.

In all experiments, ice nucleation in the epidermal layers initially occurred in the apoplast. We showed direct evidence that subsequent ice propagation across the cell wall leading to freezing of the intracellular space was visibly blocked by cold acclimation or  $\text{CaCl}_2$  treatment. Ice blockage appears to be associated with alterations of the structure and composition of cell walls through strengthening of the ice propagation barrier. Reductions in cell wall permeability, which can affect ice spread between the extra- and intracellular space, was a consistent response to cold acclimation. The decrease in cell wall permeability after cold acclimation effectively avoided ice penetration into the intracellular space, thus avoiding freezing injury. Changes in cell wall composition were closely connected with freezing resistance as a result of cold acclimation. An increased quantity of the relative crude cell wall content is consistent with other reports showing a physical augmentation of the cell wall content and association with increased freezing stress resistance. We further examined compositional changes of the cell wall during acclimation and found increased pectic polysaccharides (galacturonic acid) and elevated demethylesterification in acclimated cells. The large increase of PME activity appeared in the two-week rather than one-week cold acclimated samples which corresponded with the large increase in freezing stress resistance at two weeks. This result illustrated that the increasing freezing resistance of Japanese bunching onion more depends on the activity of PME not only the quantity of the apoplast. Furthermore, the increase of PME activity during cold acclimation with the majority of PME localized in the epidermal layer supports the observed increase in demethylesterified pectins. Collectively, these changes may provide more potential for pectin to cross-link with calcium to form a tighter gel which reduces the pore size of cell walls, and reduction in cell wall permeability. As a consequence, ice may be trapped outside the cells and freezing damage may be avoided.

Using *Allium fistulosum* as a system to study intracellular freezing avoidance is useful

since this species is easy to propagate and grow with short generation time. The species is both drought and low temperature stress resistant and the epidermal layer can be easily peeled with intact large cell size which does not autofluoresce. The results from this study indicated that freezing resistance in *Allium fistulosum* can be improved through reductions of the cell wall permeability, which is associated with pectin dynamics in response to cold acclimation.

## 6.0 FUTURE WORK AND SIGNIFICANCE

Based on this study, Japanese bunching onion (*Allium fistulosum* L.) appears to be a good model system to better understand the mechanisms of intracellular freezing stress avoidance and cold acclimation-induced alterations of the cell wall structure and composition (in particular pectins) to avoid freezing injury. However, more work on this topic is needed to further determine the specific mechanism of cell wall-regulated freezing stress resistance.

Future studies can focus on detecting the cold-induced modifications of the cell wall structures, especially pectins, in *Allium fistulosum* after cold acclimation. The dynamics of cell wall structure and composition can be tested through Synchrotron Radiation-Fourier Transform Infrared (SR-FTIR) Spectroscopy in Canadian Light Source. The pore size of cell walls will be measured, which can provide more direct evidence on the decreasing permeability of cell walls during cold acclimation.

Ca<sup>2+</sup> plays an essential role in controlling cell wall integrity, cell wall pore size and cohesion, and signal transduction as a second messenger in response to stress conditions (Hirschi, 2004). The apoplastic Ca<sup>2+</sup> can bind the contiguous un-methylesterified pectin (mainly homogalacturonan) to form the crossing-linkage which confer the rigidity of cell wall and maintain the integrity of cell wall in order to relieve the stresses (Hamann *et al.*, 2009). Consequently, the localization, quantity and pattern of calcium also can be carried out through the x-ray beamline in the Canadian Light Source, thereby better understanding how the calcium plays the role in enhancing the ability of plant to resist freezing stress. Moreover, the mechanisms on how PME activity is regulated can be tested since we have *Arabidopsis* mutants with PME1 OE (PME inhibitor overexpression), which may provide the molecular bases of the resistance to freezing stress. Therefore, the results from those tests can further confirm the roles of cell wall as a barrier in decreasing even avoiding the freezing damage.

The cell wall as a barrier may play an even wider role in stress resistance. Water loss (dehydration) is caused by extracellular freezing-induced dehydration (Yamazaki *et al.*, 2009), but also desiccation and salt stress (Zhu, 2002) and postharvest desiccation (Belincontro *et al.*, 2004). Ice penetration can be prevented by the decreasing cell wall permeability which is determined by the pore size of cell walls. There is some evidence that pectins are involved in the permeability barrier to limit the ice propagation (Jones *et al.*, 2000) because the acidic pectins

are capable of forming the pectate gel with  $\text{Ca}^{2+}$ . The acidic pectins which are un-methylated are the products of the hydrolysis of methylated pectins under the catalysis of pectin methylesterase (PME), so the pectin gelation is directly regulated by the PME activity. Therefore, PME activity in the freezing resistance is important since PME controls the cell wall permeability that can reduce and even avoid injury from ice penetration to the internal cells.

We found PME activity was highest in the epidermal layer and increased during cold acclimation in both the epidermal layer and underlying non-epidermal layers. A similar result was also found in the Solecka *et al.*'s research (2008) in which tissue resistance to blackleg fungus may have been related to the modifications of leaf cell wall structure and composition (especially pectin) in response to low temperature. PME gene expression also increases upon cold-treatment of *Arabidopsis* pollen (Lee, 2003).

PME contributes to the resistance of plants to pathogens and the plants having low PME activity are more resistant to various pathogens (McMillan *et al.*, 1993; Marty *et al.*, 1997; Boudart *et al.*, 1998; Wydra and Berl, 2006; Lionetti *et al.*, 2007; Lionetti *et al.*, 2012). Consequently, research on PME in response to freezing and pathogens should be expanded to different plant species exposed to various abiotic and biotic stresses. Based on desired PME property, breeders can select for freezing avoidance (high PME activity) or pathogen resistance (low PME activity). As a consequence, the study on PME may provide a possibility to directly select the freezing- or pathogen- resistant species. This possibility also can be examined in other stress conditions such as drought and salt stress.

In addition, we need to understand any mechanisms to resist the environmental stress not only depend on one aspect such as cell wall (mainly pectin) but also rely on other factors like plasma membrane. Actually, most of researches on freezing resistance focus on the roles of plasma membrane. Understanding the whole cellular functions in the process of freezing resistance is very critical in a comprehensive way.

## 7.0 REFERENCES

- Albersheim, P. and Anderson-Prouty, A.J. (1971) Proteins from plant cell inhibit polygalacturonase secreted by plant pathogens. *Proceedings of The National Academy of Sciences of The United States of America* **68**, 1815-1819.
- Albersheim, P. and Anderson-Prouty, A.J. (1975) Carbohydrates, proteins, cell-surfaces and biochemistry of pathogenesis. *Annual Review of Plant Physiology and Plant Molecular Biology* **26**, 31-52.
- Albersheim, P., Darvill A., Roberts K., Sederoff R., Staehelin A. (2011) Plant Cell Walls, from Chemistry to Biology. Garland Science, New York, pp 227-272.
- Alonso, A., Queiroz, C.S. and Magalhaes, A.C. (1997) Chilling stress leads to increased cell membrane rigidity in roots of coffee (*Coffea arabica* L.) seedlings. *Biochimica et Biophysica Acta* **1323**, 75-84.
- Anisimov, A.V. and Egorov, A.G. (2002) Plasmodesmata as a modulator of osmotic water fluxes in plants. *Russian Journal of Plant Physiology* **49**, 677-684.
- Anthon, G.E. and Barrett, D.M. (2006) Potential and limitations for determining lycopene in tomatoes by optical methods. *Acta Horticulturae* **724**, 243-248.
- Antikainen, M. and Griffith, M. (1997) Antifreeze protein accumulation in freezing-tolerant cereals. *Physiologia Plantarum* **99**, 423-432.
- Antikainen, M., Griffith, M., Zhang, J., Hon, W.-C, Yang, D.S.C. and Pihakaski-Maunsbach, K. (1996) Immunolocalization of antifreeze proteins in winter rye leaves, crowns, and roots by tissue printing. *Plant Physiology* **110**, 845-857.
- Aoyama, S. and Yamamoto, Y. (2007) Antioxidant activity and flavonoid content of Welsh onion (*Allium fistulosum*) and the effect of thermal treatment. *Food Science and Technology Research* **13**(1), 67-72.
- Apel, K. and Hirt, H. (2004) Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annual Review of Plant Biology* **55**, 373-399.
- Arora, R. and Palta, J.P. (1988) *In vivo* perturbation of membrane-associated calcium by freeze-thaw stress in onion bulb cells. Simulation of this perturbation in extracellular KCl and alleviation by calcium. *Plant Physiology* **87**, 622-628.

- Asahina, E. (1956) The freezing process of plant cell. *Contribution from the Institute of Low Temperature Science* **10**, 83-126.
- Ashworth, E.N. (1984) Xylem developmemt in *Prunus* flower buds and the relationship to deep supercooling. *Plant Physiology* **74**, 862-65.
- Ashworth, E.N. and Abeles, F.B. (1984) Freezing behavior of water in small pores and the possible role in the freezing of plant tissue. *Plant Physiology* **76**, 201-204.
- Ashworth, E.N. and Kieft, T.L. (1995) Ice nucleation activity associated with plants and fungi. pp. 137-162 in RE Lee, GJ Warren, and LV Gusta, (eds.) Biological ice nucleation and its applications, APS Press, St. Paul, MN.
- Ashworth, E.N. and Pearce, R.S. (2002) Extracellular freezing in leaves of freezing-sensitive species. *Planta* **214**, 798-805.
- Ashworth, E.N., Echlin, P., Pearce, R.S. and Hayes, T.L. (1988) Ice formation and tissue response in apple twigs. *Plant, Cell and Environment* **11**, 703-710.
- Baldwin, L., Domon, J.M., Klimek, J.F., Fournet, F., Sellier, H., Gillet, F., Pelloux, J., Lejeune-Hénaut, I., Carpita, N.C. and Rayon, C. (2014) Structural alteration of cell wall pectins accompanies pea development in response to cold. *Phytochemistry* **104**, 37-47.
- Baron-Epel, O., Gharyl, P.K. and Schindler, M. (1988) Pectins as mediators of wall porosity in soybean cells. *Planta* **175**, 389-395.
- Bashline, L., Lei, L., Li, S. and Gu, Y. (2014) Cell wall, cytoskeleton, and cell expansion in higher plants. *Molecular Plant* **7**, 586-600.
- Bateman, D.F. and Basham, H.G. (1976) Degradation of plant cell walls and membranes by microbial enzymes. pp. 316-355 in R Heitefuss and PH Williams (eds.), *Encyclopedia of Plant Physiology. Physiological Plant Pathology*, Springer-Verlag, Berlin.
- Bellincotro, A., De Santis, D., Botondi, R., Villa, I. and Mencarelli, F. (2004) Different postharvest dehydration rates affect quality characteristics and volatile compounds of Malvasia, Trebbiano and Sangiovese grapes for wine production. *Journal of the Science of Food and Agriculture* **84**, 1791-1800.
- Biermann, J., Stushnoff, C. and Burke, M.J. (1979) Differential thermal analysis and freezing injury in cold hardy blueberry flower buds. *Journal of the American Society for Horticultural Science* **104**, 444-449.
- Bordenave, M. and Goldberg, R. (1993) Purification and characterization of pectin

- methylesterases from mung bean hypocotyl cell walls. *Phytochemistry* **33**, 999-1003.
- Bosch, M. (2005) Pectin methylesterases and pectin dynamics in pollen tubes. *The Plant Cell* **17(12)**, 3219-3226.
- Boudart, G., Lafitte, C., Barthe, J.P., Frasez, D. and Esquerré-Tugayé, M.T. (1998) Differential elicitation of defense responses by pectic fragments in bean seedlings. *Planta* **206**, 86-94.
- Brett, C.T. and Waldron, K.W. (1996) Physiology and Biochemistry of Plant Cell Walls, 2<sup>nd</sup> ed, Chapman and Hall, London.
- Brummell, D.A., Dal Cin, V., Lurie, S., Crisosto, C.H. and Labavitch, J.M. (2004) Cell wall metabolism during the development of chilling injury in cold-stored peach fruit: association of mealiness with arrested disassembly of cell wall pectins. *Journal of Experimental Botany* **55(405)**, 2041-2052.
- Buchanan, B., Gruissem, W. and Jones, R. (2000) Biochemistry and molecular biology of plants, 1<sup>st</sup> ed, American Society of Plant Physiology.
- Buchanan, C.D., Lim, S., Salzman, R.A., Kagiampakis, I., Morishige, D.T., Weers, B.D., Klein, R.R., Pratt, L.H., Cordonnier-Pratt, M.M. and Klein, P.E. (2005) *Sorghum bicolor*'s transcriptome response to dehydration, high salinity and ABA. *Plant Molecular Biology* **58**, 699-720.
- Buckeridge, M.S., Santos, H.P. and Tiné, M.A.S. (2000) Mobilisation of storage cell wall polysaccharides in seeds. *Plant Physiology and Biochemistry* **38**, 141.
- Burchett, S., Niven, S. and Fuller, M.P. (2006) The effect of cold-acclimation on the water relations and freezing tolerance *Hordeum vulgare* L. *Cryo Letters* **27**, 295-303.
- Burke, M.J. and Lindow, S.E. (1990) Surface properties and size of the ice nucleation site in ice nucleation active bacteria considerations. *Cryobiology* **27**, 80-84.
- Burke, M.J., Gusta, L.V., Quamme, H.A., Weiser, C.J. and Li, P.H. (1976) Freezing and injury in plants. *Annual Review of Plant Physiology* **27**, 507-528.
- Burström, H.G. (1968). Calcium and plant growth. *Biological Review (Cambridge)* **43**, 287-316.
- Bush, D.S. (1993) Regulation of cytosolic calcium in plants. *Plant Physiology* **103**, 7-13.
- Bush, D.S. (1995) Calcium Regulation in Plant Cells and Its Role in Signalling. *Annual Review of Plant Physiology and Plant Molecular Biology* **46**, 95-122.
- Caffall, K.H. and Mohnen, D. (2009) The structure, function, and biosynthesis of plant cell wall pectic polysaccharides. *Carbohydrate Research* **344**, 1879-1900.



- Carpita, N.C., Sabularse, D., Montezinos, D. and Delmer, D.P. (1979) Determination of the pore size of cell walls of living plant cells. *Science* **205**, 1144-1147.
- Carpita, N.C. and Gibeaut, D.M. (1993) Structural models of primary cell walls in flowering plants: consistency of molecular structure with the physical properties of the walls during growth. *The Plant Journal* **3**, 1-30.
- Carvajal, M., Cooke, D.T. and Clarkson, D.T. (1998) The lipid bilayer and aquaporins: parallel pathways for water movement into plant cells. *Plant Growth Regulation* **25**, 89-95.
- Carvalho, C.P., Hayashi, A.H., Braga, M.R. and Nievola, C.C. (2013) Biochemical and anatomical responses related to the *in vitro* survival of the tropical bromeliad *Nidularium minutum* to low temperatures. *Plant Physiology and Biochemistry* **71**, 144-154.
- Chalker-Scott, L. (1992) Disruption of an ice-nucleation barrier in cold hardy *Azalea* buds by sublethal heat stress. *Annals of Botany* **70**, 409-418.
- Chang, C.Y., Lai, L.R. and Chang, W.H. (1995) Relationships between textural changes and the changes in linkages of pectic substances of sweet pepper during cooking processes, and the applicability of the models of interactions between pectin molecules. *Food Chemistry* **53**, 409-416.
- Chanliaud, E., Burrows, K.M., Jeronimidis, G. and Gidley, M.J. (2002) Mechanical properties of primary plant cell wall analogues. *Planta* **215**, 989-996.
- Chen, P.M., Li, P.H. and Cunningham, W.P. (1977) Ultrastructure difference in leaf cell of *Solanum* species in relation to their frost resistance. *Botanical Gazette* **138**, 267-285.
- Chinnusamy, V., Zhu, J. and Zhu, J-K. (2007) Cold stress regulation of gene expression in plants. *Trends in Plant Science* **12 (10)**, 444-451.
- Clausen, M.H., Willats, W.G.T. and Knox, J.P. (2003) Synthetic methyl hexagalacturonate hapten inhibitors of anti-homogalacturonan monoclonal antibodies LM7, JIM5 and JIM7. *Carbohydrate Research* **338**, 1797-1800.
- Clifford, S.C., Arndt, S.K., Corlett, J.E., Joshi, S., Sankhla, N., Popp, M. and Jones, H.G. (1998) The role of solute accumulation, osmotic adjustment and changes in cell wall elasticity in drought tolerance in *Ziziphus mauritiana* (Lamk.). *Journal of Experimental Botany* **49(323)**, 967-977.
- Cline, K. and Albersheim, P. (1981) Host-pathogen interactions. *Plant Physiology* **68**, 207-220.
- Close, T.J. (1997) Dehydrins: a commonality in the response of plants to dehydration and low

- temperature. *Plant Physiology* **100**, 291-296.
- Cosgrove, D.J. (2005) Growth of the plant cell wall. *Nature Reviews Molecular cell Biology* **6(11)**, 850-861.
- Dahal, K., Kane, K., Gadapati, W., Webb, E., Savitch, L.V., Singh, J., Sharma, P., Sarhan, F., Longstaffe, F.J., Grodzinski, B. and Hüner, N.P.A. (2012) The effects of phenotypic plasticity on photosynthetic performance in winter rye, winter wheat, and *Brassica napus*. *Physiologia Plantarum* **144**, 169-188.
- de Freitas, S.T., Handa, A. K., Wu, Q., Park, S. and Mitcham, J. (2012) Role of pectin methylesterases in cellular calcium distribution and blossom-end rot development in tomato fruit. *The Plant Journal* **71**, 824-835.
- de Lima, R.B., dos Santos, T.B., Vieira, L.G.E., Ferrarese, M.D.L., Ferrarese, O., Donatti, L., Boeger, M.R.T. and Petkowicz, C.L.D. (2014) Salt stress alters the cell wall polysaccharides and anatomy of coffee (*Coffea arabica* L.) leaf cells. *Carbohydrate Polymers* **112**, 686-694.
- Demarty, M., Morvan, C. and Thellier, M. (1984) Calcium and the cell wall. *Plant, Cell & Environment* **7**, 441-448.
- Dowgert, M.F. and Steponkus, P.L. (1984) Behavior of the plasma membrane of isolated protoplasts during a freeze-thaw cycle. *Plant Physiology* **75**, 1139-1151.
- Draye, M. and van Cutsem, P. (2008) Pectin methylesterases induce an abrupt increase of acidic pectin during strawberry fruit ripening. *Journal of Plant Physiology* **165**, 1152-1160.
- Duman, J.G. and Olsen, T.M. (1993) Thermal hysteresis protein-activity in bacteria, fungi and phylogenetically diverse plants. *Cryobiology* **30**, 322-328.
- Easterling, D.R. and Horton, B. (1997) Maximum and minimum temperature trends for the globe. *Science* **227**, 364-367.
- Emanuelsson, J. (1984) Root growth and calcium uptake in relation to calcium concentration. *Plant Soil* **78**, 325-334.
- Endoh, K., Kasuga, J., Arakawa, K., Ito, T. and Fujikawa, S. (2009) Cryo-scanning electron microscopic study on freezing behaviors of tissue cells in dormant buds of larch (*Larix kaempferi*). *Cryobiology* **59**, 214-222.
- Farrant, J.M. (2000) Comparison of mechanisms of desiccation tolerance among three angiosperm resurrection plants. *Plant Ecology* **151**, 29-39.

- Fenwick, K.M., Jarvis, M.C. and Apperley, D.C. (1997) Estimation of polymer rigidity in cell walls of growing and nongrowing celery collenchyma by solid-state nuclear magnetic resonance in vivo. *Plant Physiology* **115**, 587-592.
- Ferullo, J.M., Vézina, L.P., Rail, J., Laberge, S., Nadeau, P. and Castonguay, Y. (1997) Differential accumulation of two glycine-rich proteins during cold-acclimation alfalfa. *Plant Molecular Biology* **33**, 625-633.
- Ford-Lloyd, B.V. and Armstrong, S.J. (1993) Welsh onion-*Allium fistulosum* L. In G Kalloo and BO Bergh, eds., Genetic improvement of vegetable crops. Pergamon Press, London, pp 51-58.
- Foster, C.E., Martin, T.M. and Pauly, M. (2010) Comprehensive Compositional Analysis of Plant Cell Walls (Lignocellulosic biomass) Part II: Carbohydrates. *Journal of Visualized Experiments* **37**, 1837-1841.
- Fry, S.C. (1986) Cross-linking of matrix polymers in the growing cell walls of angiosperms. *Annual Review of Plant Physiology* **37**, 165-186.
- Fujikawa, S. and Kuroda, K. (2000) Cryo-scanning electron microscopic study on freezing behavior of xylem ray parenchyma cells in hardwood species. *Micron* **31**, 669-686.
- Fujikawa, S., Kasuga, J., Takata, N. and Arakawa, K. (2009) Factors related to change of deep supercooling capability in xylem parenchyma cells of trees. pp. 29-42 in Gusta, L. V., Wisniewski, M. and Tanino, K. (eds.) *Plant Cold Hardiness: From the Laboratory to the Field*, CABI Press, Wallingford, UK.
- Fujikawa, S., Kuroda, K. and Fukazawa, S. (1994) Ultrastructural study of deep supercooling of xylem ray parenchyma cells from *Styrax obassia*. *Micron* **25**, 241-252.
- Fujikawa, S., Kuroda, K., Jitsuyama, Y., Sano, Y. and Ohtani, J. (1999) Freezing behavior of xylem ray parenchyma cells in softwood species with differences in the organization of cell walls. *Protoplasma* **206**, 31-40.
- Futamura, N., Mori, H., Kouchi, H. and Shinohara, K. (2000) Male flower-specific expression of genes for polygalacturonase, pectin methylesterase and  $\beta$ -1,3-glucanase in a dioecious willow (*Salix gilgiana* Seemen). *Plant Cell Physiology* **41**, 16-26.
- Gall, H.L., Philippe, F., Domon, J.M., Gillet, F., Pelloux, J. and Rayon, C. (2015) Cell wall metabolism in response to abiotic stress. *Plants* **4**, 112-166.
- Gao, H.B., Chen, G.L., Han, L.H., and Lin, H. (2004) Calcium influence on chilling resistance of

- grafting eggplant seedlings. *Journal of Plant Nutrition* **27**, 1327-1339.
- Geitmann, A. (2010) Mechanical modeling and structural analysis of the primary plant cell wall. *Current Opinion in Plant Biology* **13**, 693-699.
- Gilmour, S.J., Artus, N.N. and Thomashow, M.F. (1992) cDNA sequence analysis and expression of two cold-regulated genes of *Arabidopsis thaliana*. *Plant Molecular Biology* **18**, 13-21.
- Goldberg, R., Morvan, C., Jauneau, A. and Jarvis, M.C. (1996) Methyl-esterification, de-esterification and gelation of pectins in the primary cell wall. *Progress in Biotechnology* **14**, 151-172.
- Gong, S.Y., Huang, G.Q., Sun, X., Li, P., Zhao, L.L., Zhang, D.J. and Li, X.B. (2011) GhAGP31, a cotton non-classical arabinogalactan protein, is involved in response to cold stress during early seedling development. *Plant Biology* **14**, 447-457.
- Graham, P.R. and Mullin, R. (1976) The determination of lethal freezing temperatures in buds and stems of deciduous azalea by a freezing curve method. *Journal of the American Society for Horticultural Science* **101**, 3-7.
- Grant, G.T., Morris, E.R., Rees, D.A., Smith, P.J.C. and Thom, D. (1973) Biological interactions between polysaccharides and divalent cations: the egg-box model. *FEBS Letters* **32**, 195-198.
- Grevsen, K. (1989) Effects of sowing dates on different varieties of Welsh onion (*Allium fistulosum* L.) under temperature coastal climate. *Acta Horticulture* **242**, 319-324.
- Griffith, M. and Brown, G.N. (1982) Cell wall deposits in winter rye *Secale cereale* L. 'Puma' during cold acclimation. *Botanical Gazette* **143**, 486-490.
- Griffith, M. and Yaish, M.W. (2004) Antifreeze proteins in overwintering plants: a tale of two activities. *Trends in Plant Sciences* **9(8)**, 399-405.
- Griffith, M., Ala, P., Yang, D.S.C., Hon, W.C. and Moffatt, B.A. (1992) Antifreeze protein produced endogenously in winter rye leaves. *Plant Physiology* **100**, 593-596.
- Griffith, M., Antikainen, M., Hon, W.C., Pihakaski-Maunsbach, K., Yu, X.M., Chun, J.U. and Yang, D.S.C. (1997) Antifreeze proteins in winter rye. *Physiologia Plantarum* **100**, 327-332.

- Griffith, M., Huner, N.P.A., Espelie, K.E. and Kolattukudy, P.E. (1985) Lipid polymers accumulate in the epidermis and mesophyll sheath cell walls during low temperature development of winter rye leaves. *Protoplasma* **125**, 53-64.
- Griffith, M., Lumb, C., Wiseman, S.B., Wisniewski, M., Johnson, R.W. and Marangoni, A.G. (2005) Antifreeze proteins modify the freezing process in plants. *Plant Physiology* **138**, 330-340.
- Gu, L., Hanson, P.J., Post, W.M., Kaiser, D.P., Yang, B., Nemani, R., Pallardy, S.G. and Meyers, T. (2008) The 2007 eastern US spring freeze: Increased cold damage in a warming world. *BioScience* **58**, 253-262.
- Guglielmino, N., Liberman, M., Catesson, A.M., Marek, A., Prat, R., Mutaftchiev, S. and Goldberg, R. (1997b) Pectin methylesterases from poplar cambium and inner bark: Localization, properties and seasonal changes. *Planta* **202**, 70-75.
- Guglielmino, N., Liberman, M., Jauneau, A., Vian, B., Catesson, A.M. and Goldberg, R. (1997a) Pectin immunolocalization and calcium visualization in differentiating derivatives from poplar cambium. *Protoplasma* **199**, 151-160.
- Guillemin, A., Guillon, F., Degraeve, P., Rondeau, C., Devaux, M.F., Huber, F., Badel, E., Saurel, R., Lahaye, M. (2008) Firming of fruit tissues by vacuum-infusion of pectin methylesterase: visualisation of enzyme action. *Food Chemistry* **109**, 368-378.
- Gusta, L.V. and Wisniewski, M. (2013) Understanding plant cold hardiness: an opinion. *Physiologia Plantarum* **147** (1), 4-14.
- Gusta, L.V., Burke, M.J. and Kapoor, A.C. (1975) Determination of unfrozen water in winter cereals at subfreezing temperatures. *Plant Physiology* **56**, 707-709.
- Gusta, L.V., Wisniewski, M., Nesbitt, N.T. and Gusta, M.L. (2004) The effect of water, sugars, and proteins on the pattern of ice nucleation and propagation in acclimated and nonacclimated canola leaves. *Plant Physiology* **135**, 1642-1653.
- Gusta, L.V., Wisniewski, M.E. and Trischuk, R.G. (2009) Patterns of freezing in plants: the influence of species, environment and experimental conditions, pp. 214-225 in Gusta, L., Wisniewski, M. and Tanino, K. (eds.) *Plant Cold Hardiness: From the Laboratory to the Field*. CABI, Oxford, UK.
- Guy, C.L. (1990) Cold acclimation and freezing stress tolerance: role of protein metabolism. *Annual Review of Plant Physiology and Plant Molecular Biology* **41**, 187-223.

- Guy, C.L. (2003) Freezing tolerance of plants: current understanding and selected emerging concepts. *Canadian Journal of Botany* **81**, 1216-1223.
- Ha, M.A., Apperley, D.C. and Jarvis, M. (1997) Molecular rigidity in dry and hydrated onion cell wall. *Plant Physiology* **115**, 593-598.
- Hacker, J., Ladinig, U., Wagner, J. and Neuner, G. (2011) Inflorescences of alpine cushion plants freeze autonomously and may survive subzero temperatures by supercooling. *Plant Science* **180**, 149-156.
- Hahn, M.G., Bucheli, P., Cervone, F., Doares, S.H., O'Neill, R.A., Darvill, A. and Albersheim, P. (1989) Roles of cell wall constituents in plant-pathogen interactions. pp. 131-181 in Nester, E. and Kosuge, T. (eds.) *Plant-microbe interactions*, Vol 3, Mc Graw Hill, New York.
- Hamann, T., Bennett, M., Mansfield, J. and Somerville, C. (2009) Identification of cell-wall stress as a hexose-dependent and osmosensitive regulator of plant responses. *Plant Journal* **57**, 1015-026.
- Han, Y., Wang, W., Sun, J., Ding, M., Zhao, R., Deng, S., Wang, F., Hu, Y., Wang, Y., Lu, Y., Du, L., Hu, Z., Diekmann, H., Shen, X., Polle, A. and Chen, S. (2013) *Populus euphratica* XTH overexpression enhances salinity tolerance by the development of leaf succulence in transgenic tobacco plants. *Journal of Experimental Botany* **64**, 4225-4238.
- Hanelt, P. (1990) Taxonomy, evolution, and history. pp. 1-26 in Rabinowitch, H. D. and Brewster, J. L. (eds.), *Onions and Allied Crops. Vol. I. Botany, Physiology, and Genetics*. CRC Press.
- Hanin, M., Brini, F., Ebel, C.H., Toda, Y., Takeda, S. and Masmoudi, K. (2011) Plant dehydrins and stress tolerance. versatile proteins for complex mechanisms. *Plant Signaling & Behavior* **6**, 1503-1509.
- Hara, M. (2010) The multifunctionality of dehydrins. an overview. *Plant Signaling & Behavior* **5**, 503-508.
- Harholt, J., Suttangkakul, A. and Scheller H.V. (2010) Biosynthesis of pectin. *Plant Physiology* **153**, 384-395.
- Harper, L. and Freeling, M. (1996) Interactions of *liguleless1* and *liguleless2* function during ligule induction in maize. *Genetics* **144**, 1871-1882.

- Hart, D.A. and Kindel, P.K. (1970) Isolation and partial characterization of apiogalacturonans from the cell wall of *Lemna minor*. *Biochemical Journal* **116**(4), 569-579.
- Hartung, W., Schiller, P. and Dietz, K.J. (1998) Physiology of poikilohydric plants. *Progress in Botany* **59**, 299-327.
- Hepler, P.K. (2005) Calcium: A central regulator of plant growth and development. *The Plant Cell* **17**, 2142-2155.
- Hepler, P.K. and Wayne, R.O. (1985) Calcium and Plant Development, *Annual Review of Plant Physiology and Plant Molecular Biology* **36**, 397-439.
- Hincha, D.K. and Hagemann, M. (2004) Stabilization of model membranes during drying by compatible solutes involved in the stress tolerance of plants and microorganisms. *Biochemistry Journal* **383**, 277-283.
- Hirschi, K.D. (2004) The calcium conundrum. Both versatile nutrient and specific signal. *Plant Physiology* **136**, 2438-2442.
- Hon, W-C., Griffith, M., Mlynarz, A., Kwok, Y.C. and Yang, D.S.C. (1995) Antifreeze proteins in winter rye are similar to pathogenesis-related proteins. *Plant Physiology* **109**, 879-889.
- Hon, W-C., Griffith, M., Chong, P. and Yang, D.S.C. (1994) Extraction and isolation of antifreeze proteins from winter rye (*Secale cereale* L.) leaves. *Plant Physiology* **104**, 971-980.
- Hu, Z-M., Wang, R.R., Larson, S.R., Palazzo, A., Asay, K.H. and Chatterton, J. (2001) Selection responses for molecular markers associated with anthocyanin coloration and low temperature growth traits in crested wheatgrasses. *Canadian Journal of Plant Science* **81**, 665-671.
- Huang, T. and Duman, J.G. (2002) Cloning and characterization of a thermal hysteresis/antifreeze protein with DNA-binding activity from winter bittersweet nightshade, *Solanum dulcamara*. *Plant Molecular Biology* **48**, 339-350.
- Hüner, N.P.A., Palta J.P., Li P.H. and Carter J.V. (1981) Anatomical changes in leaves of Puma rye in response to growth at cold-hardening temperatures. *Botanical Gazette* **142**, 55-62.
- Hüner, N.P.A., Öquist, G., Hurry, V.M., Krol, M., Falk, S. and Griffith, M. (1993) Photosynthesis of photo inhibition and low temperature acclimation in cold tolerant plants. *Photosynthesis Research* **37**, 19-39.

- Hurry, V.M., Malmberg, G., Gardeström, P. and Öquist, G. (1994) Effects of a short-term shift to low temperature and of long-term cold hardening on photosynthesis and ribulose 1,5-bisphosphate carboxylase/oxygenase and sucrose phosphate synthase activity in leaves of winter rye (*Secale cereale* L.). *Plant Physiology* **106**, 983-990.
- Iljin, W.S. (1933) Über den Kältetod der Pflanzen und seine Ursachen. *Protoplasma* **20**, 105-124.
- Imin, N., Kerim, T., Rolfe, B.G. and Weinman, J.J. (2004) Effect of early cold stress on the maturation of rice anthers. *Proteomics* **4**, 1873-1882.
- Inden, H. and Asahira, T. (1990) Japanese bunching onion (*Allium fistulosum* L.). pp. 159-178 in Brewster, J. L. and Rabinowitch, H. D. (eds) *Onions and Allied Crops, Vol. III. Biochemistry, Food Science, and Minor Crops*. CRC Press, Boca Raton, Florida.
- Ishii, T. (1997) Structure and functions of feruloylated polysaccharides. *Plant Science* **127**, 111-127.
- Ishikawa, M. and Sakai, A. (1981) Freezing avoidance mechanisms by supercooling in some *Rhododendron* flower buds with reference to water relations. *Plant Cell Physiology* **22**, 953-967.
- Ishikawa, M. and Sakai, A. (1982) Characteristics of freezing avoidance with freezing tolerance: a demonstration of extraorgan freezing. pp. 325-340 in Li, P. H. and Sakai, A. (eds.) *Plant Cold Hardiness and Freezing Stress*, Vol. 2, Academic Press, New York.
- Ishikawa, M., Ishikawa, M., Toyomasu, T., Aoki, T. and Price, W.S. (2015) Ice nucleation activity in various tissues of *Rhododendron* flower buds: their relevance to extraorgan freezing. *Frontiers in Plant Science* **6**, 149-161.
- Jarvis, M.C. (1984) Structure and properties of pectin gels in plant cell walls. *Plant Cell and Environment* **7**, 153-164.
- Jarvis, M.C. and McCann, C. (2000) Macromolecular biophysics of the plant cell wall: concepts and methodology. *Plant Physiology and Biochemistry* **38**, 1-13.
- Jiang, Y.W., Yao, Y. and Wang, Y. (2012) Physiological response, cell wall components, and gene expression of Switchgrass under short-term drought stress and recovery. *Crop Science* **52**, 2718-2727.
- Johansson, N. and Krull, E. (1970) Ice formation, cell contraction, and frost killing of wheat plants. *National Sweden Institute for Plant Protection Contributions* **14(131)**, 343-362.



- Jolie, R., Duvetter, T., Van Loey, A.M. and Hendrickx, M.E. (2010) Pectin methylesterase and its proteinaceous inhibitor: a review. *Carbohydrate Research* **345**, 2583-2595.
- Jones, K.S., McKersie, B.D. and Paroschy, J. (2000) Prevention of ice propagation by permeability barriers in bud axes of *Vitis vinifera*. *Canadian Journal of Botany* **78**, 3-9.
- Jones, L. and McQueen-Mason, S. (2004) A role for expansins in dehydration and rehydration of the resurrection plant *Craterostigma plantagineum*. *FEBS Lett* **559**, 61-65.
- Jones, L., Milne, J. L., Ashford, D., McCann, M.C. and McQueen-Mason, S.J. (2005) A conserved functional role of pectic polymers in stomatal guard cells from a range of plant species. *Planta* **221**, 255-264.
- Jones, L., Milne, J.L., Ashford, D. and McQueen-Mason, S.J. (2003) Cell wall arabinan is essential for guard cell function. *Proceedings of the National Academy of Sciences of the United States of America* **100**, 11783-11788.
- Kagan-Zur, V., Tieman, D.M., Marlow, S.J. and Handa, A.K. (1995) Differential regulation of polygalacturonase and pectin methylesterase gene expression during and after heat stress in ripening tomato (*Lycopersicon esculentum* Mill.) fruits. *Plant Molecular Biology* **29**, 1101-1110.
- Kalcsits, L., Silim, S. and Tanino, K. (2009). Temperature influence on dormancy induction in woody plants. pp. 108-118 in Gusta, L. V., Wisniewski, M. and Tanino, K. (eds) *Plant Cold Hardiness from the Laboratory to the Field*. CABI Press, Wallingford, UK.
- Kasuga, J., Fukushi, Y., Kuwabara, C., Wang, D., Nishioka, A., Fujikawa, E., Arakawa, K. and Fujikawa, S. (2010) Analysis of supercooling-facilitating (anti-ice nucleation) activity of flavonol glycosides. *Cryobiology* **60**, 240-243.
- Keegstra, K. (2010) Plant cell walls. *Plant Physiology* **154**, 483-486.
- Knight, H. (2000). Calcium signaling during abiotic stress in plants. *International Review of Cytology* **195**, 269-325.
- Knight, H., Trewavas, A.J. and Knight, M.R. (1996) Cold calcium signaling in Arabidopsis involves two cellular pools and a change in calcium signature after acclimation. *The Plant Cell* **8**, 489-503.
- Knox, J.P. (2008) Revealing the structural and functional diversity of plant cell walls. *Current Opinion in Plant Biology* **11**, 308-313.

- Knox, J.P., Linstead, P.J., King, J., Cooper, C., Roberts, K. (1990) Pectin esterification is spatially regulated both within cell walls and between developing tissues of root apices. *Planta* **181**, 512-521.
- Kobayashi, S., Kutsuna, N., Tanino, K.K., Uemura, M. and Kawamura Y. (2014) Confocal cryomicroscopic analysis and cryodynamics of endoplasmic reticulum in herbaceous plant cells. *Environmental and Experimental Botany* **106**, 44-51.
- Kolaksazov, M., Laporte, F., Ananieva, K., Dobrev, P., Herzog, M. and Ananiev, E.D. (2013) Effect of chilling and freezing stress on jasmonate content in *Arabis Alpine*. *Bulgarian Journal of Agricultural Science* **19**, 15-17.
- Kolattukudy, P.E. (1980) Biopolyester membranes of plants: cutin and suberin. *Science* **208**, 990-1000.
- Kołota, E., Adamczewska-Sowińska, K. and Uklańska-Pusz, C. (2013) Yielding and nutritional value of Japanese bunching onion in relation to the date of planting and type of flat cover. *Folia Horticulturae* **25(2)**, 93-100.
- Körner, C. (2003) *Alpine Plant Life: Functional Plant Ecology of High Mountain Ecosystems*. Springer-Verlag, Heidelberg.
- Kosová, K., Vitámvás, P., and Právšil, I.T. (2007). The role of dehydrins in plant response to cold. *Biologia Plantarum* **51**, 601–617.
- Kourounioti, R.L.A., Band, L.R., Fozard, J.,A., Hampstead, A., Lovrics, A., Moyroud, E., Vignolini, S., King, J.R., Jensen, O.E. and Glover, B.J. (2013) Buckling as an origin of ordered cuticular patterns in flower petals. *Journal of the Royal Society Interface* **10(80)**.
- Kozłowski, T.T. (2002) Acclimation and adaptive responses of woody plants to environmental stresses. *The Botanical Review* **68**, 270-334.
- Kozłowski, T.T., Pallardy, S.G. (1997) *Physiology of woody plants*. Academic Press, San Diego, CA.
- Kubacka-Zębalska, M. and Kacperska, A. (1999) Low temperature-induced modifications of cell wall content and polysaccharide composition in leaves of winter oilseed rape (*Brassica napus* L. var. *oleifera* L.). *Plant Science* **148**, 59-67.
- Kuo, J., O'Brien, T.P. and Canny, M.J. (1974) Pit-field distribution, plasmodesmatal frequency, and assimilate flux in the mestome sheath cells of wheat leaves. *Planta* **121**, 97-118.
- Kuroda, K., Kasuga, J., Arakawa, K. and Fujikawa, S. (2003) Xylem ray parenchyma cells in

- boreal hardwood species respond to subfreezing temperatures by deep supercooling that is accompanied by incomplete desiccation. *Plant Physiology* **131**, 736-744.
- Kuwabara, C., Wang, D., Endoh, K., Fukushi, Y., Arakawa, K. and Fujikawa, S. (2013) Analysis of supercooling activity of tannin-related polyphenols. *Cryobiology* **67**, 40-49.
- Lakowicz, J.R. (2006) Principles of Fluorescence Spectroscopy, 3<sup>rd</sup> ed., Springer, New York, USA.
- Laluk, K. and Mengiste, T. (2010). Necrotroph attacks on plants: wanton destruction or covert extortion? *The Arabidopsis Book* **8**, e0136.
- Lampert, D.T.A. (1967) Hydroxyproline-O-glycosidic linkage of the plant cell wall glycoprotein extensin. *Nature* **216**, 1322-1324.
- Lawrence, J.R., Wolfaardt, G.M. and Korber, D.R. (1994) Determination of diffusion coefficients in biofilms by confocal laser microscopy. *Applied and Environmental Microbiology* **60**, 1166-1173.
- Lazić, B., Todorović, V., Dardić, M. and Paroussi, G. (2002) Effect of production method on earliness and yield of *Allium fistulosum* L. *Acta Horticulture* **579**, 359-362.
- Lee, J.Y. (2003) Use of serial analysis of gene expression technology to reveal changes in gene expression in Arabidopsis pollen undergoing cold stress. *Plant Physiology* **132**, 517-529.
- Lee, Y.K., Derbyshire, P., Knox, J. P. and Hvoslef-Eide, A.K. (2008) Sequential cell wall transformaitons in response to the induction of a pedicel abscission event in *Euphorbia pulcherrima* (poinsettia). *The Plant Journal* **54**, 993-1003.
- Leucci, M.R. , Lenucci, M.S., Piro, G. and Dalessandro, G. (2008) Water stress and cell wall polysaccharides in the apical root zone of wheat cultivars varying in drought tolerance. *Journal of Plant Physiology* **165**, 1168-1180.
- Levitt, J. (1972) Responses of Plants to Environmental Stresses. Academic Press, New York.
- Levitt, J. (1980) Responses of Plants to Environmental Stresses, 2<sup>nd</sup> ed, Academic Press, New York.
- Levitt, J. and Scarth, G. W. (1936) Frost-hardening studies with living cells. II. Permeability in relation to frost resistance and the seasonal cycle. *Canadian Journal of Research* **14**, 285-305.
- Li, Y., Jones L., McQueen-Mason S. (2003) Expansins and cell growth. *Current Opinion of Plant Biology* **6**, 603-610

- Liesche, J., Martens, H.J. and Schulz, A. (2011) Symplasmic transport and phloem loading in gymnosperm leaves. *Protoplasma* **248**, 181-190.
- Liners, F., Letesson, J.J., Didembourg, C. and Van Cutsem, P. (1989) Monoclonal antibodies against pectin: recognition of a conformation induced by calcium. *Plant Physiology* **91**, 1419-1424.
- Lionetti, V., Cervone, F. and Bellincampi, D. (2012) Methyl esterification of pectin plays a role during plant-pathogen interactions and affects plant resistance to diseases. *Journal of Plant Physiology* **169**, 1623-1630.
- Lionetti, V., Raiola, A., Camardella, L., Giovane, A., Obel, N., Pauly, M., Favaron, F., Cervone, F. and Bellincampi, D. (2007) Overexpression of pectin methylesterase inhibitors in *Arabidopsis* restricts fungal infection by *Botrytis cinerea*. *Plant Physiology* **143**, 1871-1880.
- Lipp, C.C., Goldstein, G., Meinzer, F.C. and Niemczura, W. (1994) Freezing tolerance and avoidance in high-elevation Hawaiian plants. *Plant, Cell and Environment* **17**, 1035-1044.
- Liu, H., Chen, F., Yang, H., Yao, Y., Gong, X., Xin, Y. and Ding, C. (2009) Effect of calcium treatment on nanostructure of chelate-soluble pectin and physicochemical and textural properties of apricot fruits. *Food Research International* **42**, 1131-1140.
- Livingston, D.P. and Henson, C.A. (1998) Apoplastic sugars, fructans, fructan exohydrolase and invertase in winter oat: responses to second-phase cold-acclimation. *Plant Physiology* **116**, 403-408.
- Livingston, D.P., Hinch, D.K., Heyer, A.G. (2009) Fructan and its relationship to abiotic stress tolerance in plants. *Cellular and Molecular Life Sciences* **66**, 2007-2023.
- Lodish, H., Berk, A., Zipursky, S.L., Matsudaira, P., Baltimore, D., Darnell, J. (2000) Molecular Cell Biology. In Freeman, W. H. *The Dynamic Plant Cell Wall*, 4<sup>th</sup> ed, New York.
- Ludwig, A.A., Romeis, T. and Jones, J.D.G. (2004) CDPK-mediated signaling pathways: specificity and cross-talk. *Journal of Experimental Botany* **55**, 181-188.
- Maathuis, F.J.M. (2009) Physiological functions of mineral macronutrients. *Current Opinion in Plant Biology* **12**, 250-258.
- Mahajan, S. and Tuteja, N. (2005) Cold, salinity and drought stresses: An overview. *Archives of Biochemistry and Biophysics* **444**, 139-158.

- Mahouachi, J., Socorro, A.R. and Talon, M. (2006) Responses of papaya seedlings (*Carica papaya* L.) to water stress and re-hydration: growth, photosynthesis and mineral nutrient imbalance. *Plant Soil* **281**, 137-146.
- Mankarios, A.T., Hall, M. A., Jarvis, M.C., Threlfall, D.R. and Friend, J. (1980) Cell wall polysaccharides from onions. *Phytochemistry* **19**, 1731-1733.
- Marentes, E., Griffith, M., Mlynarz, A. and Brush, R.A. (1993) Proteins accumulate in the apoplast of winter rye leaves during cold acclimation. *Physiologia Plantarum* **87**, 499-507.
- Marty, P., Jouan, B., Bertheau, Y., Vian, B. and Goldberg, R. (1997) Charge density in stem cell walls of *Solanum tuberosum* genotypes and susceptibility to blackleg. *Phytochemistry* **44**, 1435-1441.
- Matoh, T. and Kobayashi, M. (1998) Boron and calcium, essential inorganic constituents of pectic polysaccharides in higher plant cell walls. *Journal of Plant Research* **111**, 179-190.
- Matsuura, Y., Matsubara, K. and Fuchigami, M. (2000) Molecular composition of onion pectic acid. *Food Chemistry and Toxicology* **65**, 1160-1163.
- Mazur, P. (1963) Kinetics of water loss from cells at subzero temperatures and the likelihood of intracellular freezing. *The Journal of General Physiology* **47**, 347-369.
- Mazur, P. (1969) Freezing injury in plants. *Annual Review of Plant Physiology* **20**, 419-448.
- Mazzitelli, L., Hancock, R.D., Haupt, S., Walker, P.G., Pont, S.D.A., McNicol, J., Cardle, L., Morris, J., Viola, R., Brennan, R., Hedley, P.E. and Taylor, M.A. (2007) Co-ordinated gene expression during phases of dormancy release in raspberry (*Rubus idaeus* L.) buds. *Journal of Experiment Botany* **58**, 1035-1045.
- McAinsh, M.R. and Pittman, J.K. (2009) Shaping the calcium signature. *New Phytologist* **181**, 275-294.
- McCann, M.C., Wells, B. and Roberts, K. (1990) Direct visualization of cross-links in the primary plant cell wall. *Journal of Cell Science* **96**, 323-334.
- McCann, M.C., Hammouri, M., Wilson, R., Belton, P. and Roberts, K. (1992) Fourier transform infrared microspectroscopy is a new way to look at plant cell walls. *Plant Physiology* **100**, 1940-1947.

- McDougall, G.J., Morrison, I.M., Stewart, D. and Hillman, J.R. (1996) Plant cell wall as dietary fibre: range, structure, processing and function. *Journal of the Science of Food and Agriculture* **70**, 133-150.
- McMillan, G.P., Hedley, D., Fyffe, L. and Pérombelon, M.C.M. (1993) Potato resistance to soft-rot erwinias is related to cell wall pectin esterification. *Physiological and Molecular Plant Pathology* **42**, 279-289.
- McNeil, M., Darvill, A.G., Fry, S.C. and Albersheim, P. (1984) Structure and function of the primary cell walls of plants. *Annual Review of Biochemistry* **53**, 625-663.
- Medvedev, S.S. (1997) Polarity and Embryogeny in Plants, in T.B. Batygina (ed.), *Embryology of Flowering Plants, Terminology and Concepts. Vol. 2. The Seed, World and Family*, St.-Petersburg, pp. 594–601 (in Russian).
- Medvedev, S.S. (2005) Calcium signaling system in plants. *Russian Journal of Plant Physiology* **52**, 249-270.
- Medvedev, S.S. and Markova, I.V. (1998) The Cytoskeleton and Plant Polarity. *Fiziologiya Rastenii* (Moscow) **45**, 185-197 (Russian Journal of Plant Physiology, English-language Translation).
- Meier, H. and Reid, J.S.G. (1981) Reserve polysaccharides other than starch in higher plants. pp. 418-471 in Tanner, W. and Loewus, F. A. (eds.), *Encyclopedia of Plant Physiology*, Springer, Berlin.
- Micheli, F. (2001) Pectin methylesterases: Cell wall enzymes with important roles in plant physiology. *Trends in Plant Science* **6**, 414-419.
- Micheli, F., Holliger, C., Goldberg, R. and Richard, L. (1998) Characterization of the pectin methylesterase-like gene AtPME3: a new member of a gene family comprising at least 12 genes in *Arabidopsis thaliana*. *Gene* **220**, 13-20.
- Miedes, E., Suslov, D., Vandenbussche, F., Kenobi, K., Ivakov, A., Straeten, D.V.D., Lorences, E.P., Mellerowicz, E.J., Verbelen, J.P. and Vissenberg K. (2013) Xyloglucan endotransglucosylase/hydrolase (XTH) overexpression affects growth and cell wall mechanics in etiolated *Arabidopsis* hypocotyls. *Journal of Experimental Botany* **64**, 2481-2497.
- Minorsky, P.V. and Spanswick, R.M. (1989) Electrophysiological evidence for a role for calcium in temperature sensing by roots of cucumber seedlings. *Plant, Cell & Environment* **12**,

137-143.

- Moellering, E.R., Muthan, B. and Benning, C. (2010) Freezing tolerance in plants requires lipid remodeling at the outer chloroplast membrane. *Science* **330**, 226-228.
- Mohnen, D. (2008) Pectin structure and biosynthesis. *Current Opinion in Plant Biology* **11**(3), 266-277.
- Monroy, A.F. and Dhindsa, R.S. (1995) Low temperature signal transduction: The induction of cold acclimation-specific genes of alfalfa by calcium at 25°C. *The Plant Cell* **7**, 321-331.
- Monroy, A.F., Sarhan, F., and Dhindsa, R.S. (1993) Cold-induced changes in freezing tolerance, protein phosphorylation, and gene expression: Evidence for a role of calcium. *Plant Physiology* **102**, 1227-1235.
- Moore, E.B. and Molinero, V. (2011) Structural transformation in supercooled water controls the crystallization rate of ice. *Nature* **479**, 506-508.
- Moore, J.P., Farrant, J.M. and Driouich, A. (2008a) A role for pectin-associated arabinans in maintaining the flexibility of the plant cell wall during water deficit stress. *Plant Signal and Behavior* **3**, 102-104.
- Moore, J.P., Vitré –Gibouinb, M., Farrantc, J.M. and Driouich, A. (2008b) Adaptations of higher plant cell walls to water loss: drought vs desiccation. *Physiologia Plantarum* **134**, 237-245.
- Moore, P.J. and Staehelin, L.A. (1988) Immunogold localization of the cell-wall-matrix polysaccharides rhamnogalacturonan I and xyloglucan during cell expansion and cytokinesis in *Trifolium pretense* L.; implication for secretory pathways. *Planta* **174**, 433-445.
- Muldrew, K., Acker, P. J., Elliot, A.J. and McGann, E.L. (2004) The water to ice transition: implications for living cells, pp 79-101 in Fuller, J. B., Lane, N. and Benson, E. E. (eds.), *Life in the Frozen State*, CRC Press, Boca Raton, Florida.
- Müller, K., Levesque-Tremblay, G., Bartels, S., Weibrecht, K., Wormit, A., Usadel, B., Haughn, G. and Kermode, A.R. (2013) Demethylesterification of cell wall pectins in *Arabidopsis thaliana* plays a role in seed germination. *Plant Physiology* **161**, 305-316.
- Murai, M. and Yoshida, S. (1998) Evidence for the cell wall involvement in temporal changes in freezing tolerance of Jerusalem Artichoke (*Helianthus tuberosus* L.) tubers during cold acclimation. *Plant and Cell Physiology* **39**, 97-105.

- Netzer, D., Rabinowitch, H.D. and Weintal, C. (1985) Greenhouse technique to evaluate onion resistance to pink root. *Euphytica* **34**, 385-391.
- Ng, A., Smith, A.C. and Waldron, K.W. (1998) Effect of tissue type and variety on cell wall chemistry of onion (*Allium cepa* L.). *Food Chemistry* **63**(1), 17-24.
- Ng, C.K.-Y., McAinsh, M.R., Gray, J.E., Hunt, L., Leckie, C.P., Mills, L. and Hetherington, A.M. (2001) Calcium-based signalling systems in guard cells. *New Phytologist* **151**, 109-120.
- Noh, S.A., Park, S.H., Huh, G. H., Paek, K-H., Shin, J.S. and Bae, J.M. (2009) Growth retardation and differential regulation of expansin genes in chilling-stressed sweet potato. *Plant Biotechnology Reports* **3**, 75-85.
- O'Neill, M.A., Albersheim, P. and Darvill, A., (1990) The pectic polysaccharides of primary cell walls. pp. 415-441 in Dey, D.M. (eds.), *Methods in Plant Biochemistry*, Academic Press, London.
- O'Neil, M.A. and York, W.S. (2003) The composition and structure of plant primary cell walls. In *The Plant Cell Wall*, Rose, J.K.C. (ed.), pp: 1-54. CRC Press, Oxford.
- Ober, E.S. and Sharp, R.E. (1994) Proline accumulation in maize (*Zea mays* L.) primary roots at low water potentials I. Requirement for increased levels of abscisic acid. *Plant Physiology* **105**, 981-987.
- Olien, C.R. (1961) A method of studying stresses occurring in plant tissue during freezing. *Crop Science* **1**, 26-28.
- Olien, C.R. and Clark, J.L. (1993) Changes in soluble carbohydrate composition of barley, wheat, and rye during winter. *Agronomy Journal* **85**, 21-29.
- Olien, C.R. and Lester, G.E. (1985) Freeze-induced changes in soluble carbohydrates of rye. *Crop Science* **25**, 288-290.
- Olien, C.R. and Marchetti, B.L. (1976) Recovery of hardened barley from winter injuries. *Crop Science* **16**, 201-204.
- Olien, C.R. and Smith, M.N. (1977) Adhesions in relation to freeze stress. *Plant Physiology* **60**, 499-503.
- Omran, A.O., Atkins, I.M. and Gilmore, E.C. (1968) Heritability of cold hardness in flax (*Linum usitatissimum* L.). *Crop Science* **8**, 716-719.



- Ortbauer, M. (2013) Abiotic stress adaptation: protein folding stability and dynamics. In: Abiotic stress-plant responses and applications in agriculture, pp. 3-25, Vahdati, K. and Leslie, C. (eds.). InTech, Rijeka, Croatia.
- Palta, J.P. (1990) Stress interactions at the cellular and membrane levels. *HortScience* **25**, 1377-1381.
- Palta, J.P. (1996) Role of calcium in plant responses to stresses: Linking basic research to the solution of practical problems. *HortScience* **31**, 51-57.
- Palta, J.P. and Li, P.H. (1980) Alterations in membrane transport properties by freezing injury in herbaceous plants. Evidence against rupture theory. *Physiologia Plantarum* **50**, 169-175.
- Palta, J.P., Levitt, J. and Stadelmann, E.J. (1977) Freezing injury in onion bulb cells: I. Evaluation of the conductivity method and analysis of ion and sugar efflux from injured cells. *Journal of Plant Physiology* **60**, 398-401.
- Parre, E. and Geitmann, A. (2005) Pectin and the role of the physical properties of the cell wall in pollen tube growth of *Solanum chacoense*. *Planta* **220**, 582-592.
- Pearce, R.S. (1988) Extracellular ice formation and cell shape in frost-stressed cereal leaves: a low-temperature scanning-electron-microscopy study. *Planta* **175**, 313-324.
- Pearce, R.S. (2001) Plant freezing and damage. *Annals of Botany* **87**, 417-424.
- Pearce, R.S. and Ashworth, E.N. (1992) Cell shape and localisation of ice in leaves of overwintering wheat during frost stress in the field. *Planta* **188**, 324-331.
- Pearce, R.S. and Fuller, M.P. (2001) Freezing of barley (*Hordeum*) studied by infrared video thermography. *Plant Physiology* **125**, 227-240.
- Pearce, R.S. and Willison, J.H. (1985) Wheat tissue freeze-etched during exposure to extracellular freezing: distribution of ice. *Planta* **163(3)**, 295-303.
- Peaucelle, A., Braybrook, S.A. and Höfte, H. (2012) Cell wall mechanics and growth control in plants: the role of pectins revisited. *Frontiers in Plant Physiology* **3**, 121.
- Pelloux, J., Rusterucci, C. and Mellerowicz, E.J. (2007) New insights into pectin methyl-esterase structure and function. *Trends in Plant Science* **12**, 267-277.
- Percival, G.C. and Barnes, S. (2008) Calcium-induced freezing and salinity tolerance in evergreen oak and apple cv. 'Golden Crow'. *Arboriculture and Urban Forestry* **34**, 191-199.

- Peterson, C.A. and Cholewa, E. (1998) Structural modifications of the apoplast and their potential impact on ion uptake. *Zeitschrift für Pflanzenernährung und Bodenkunde* **161**, 521-531.
- Polisensky, D.H. and Braam, J. (1996) Cold shock regulation of the *Arabidopsis* TCH genes and the effects of modulating intracellular calcium levels. *Plant Physiology* **111**, 1271-1279.
- Preston, J.C. and Sandve, S.R. (2013) Adaptation to seasonality and the winter freeze. *Frontiers in Plant Science* **4**, 167-185.
- Quamme, H.A. (1974) An exothermic process involved in the freezing injury to flower buds of several *Prunus* species. *Journal of the American Society for Horticultural Science* **99**, 315-318.
- Quamme, H.A. (1985) Avoidance of freezing injury in woody plants by deep supercooling. *Acta Horticulture* **168**, 11-27.
- Raese, J.T. (1997) Cold tolerance, yield and fruit quality of d'Anjou pears influenced by nitrogen fertilizer rates and time of application. *Journal of Plant Nutrition* **20**, 1007-1025.
- Rajashekar, C.B. and Burke, M.J. (1996) Freezing characteristics of rigid plant tissues (development of cell tension during extracellular freezing). *Plant Physiology* **111**, 597-603.
- Rajashekar, C.B. and Lafta, A. (1996) Cell wall changes and cell tension in response to cold acclimation and exogenous abscisic acid in leaves and cell cultures. *Plant Physiology* **111**, 605-612.
- Raulston, J.C. and Tripp, K.E. (1994) Exploring the complexities of plant hardiness. *Arnoldia* **54(3)**, 22-31.
- Raven, J.A. (1997) Multiple origins of plasmodesmata. *European Journal of Phycology* **32**, 95-101.
- Raymond, J.A. and DeVries, A.L. (1977) Adsorption inhibition as a mechanism of freezing resistance in polar fishes. *Proceedings of the National Academy of Sciences of the United States of America* **74**, 2589-2593.
- Ren, C. and Kermode, A.R. (2000) An increase in pectin methyl esterase activity accompanies dormancy breakage and germination of yellow cedar seeds. *Plant Physiology* **124**, 231-242.
- Richard, L., Qin, L.X. and Goldberg, R. (1996) Clustered genes within the genome of

- Arabidopsis thaliana* encoding pectin methylesterase-like enzymes. *Gene* **170**, 207-211.
- Ridley, B.L., O'Neil, M.A. and Mohnen, D. (2001) Pectins: structure, biosynthesis, and oligogalacturonide-related signaling. *Phytochemistry* **57**, 929-967.
- Rinne, P.L.H., Welling, A., Vahala, J., Ripel, L., Ruonala, R., Kangasjärvi, J. and van der Schoot, C. (2011) Chilling of dormant buds hyper induces *FLOWERING LOCUS T* and recruits GA-inducible 1,3- $\beta$ -glucanases to reopen signal conduits and release dormancy in *Populus*. *The Plant Cell* **23**, 130-146.
- Roberts, D.M. and Harmon, A.C. (1992) Calcium-modulated proteins: Targets of intracellular calcium signals in higher plants. *Annual Review of Plant Physiology and Plant Molecular Biology* **43**, 375-414.
- Rose, J.K.C. (2003) The plant cell wall. CRC Press, Oxford.
- Ruelland, E., Vaultier, M.N., Zachowski, A. and Hurry, V. (2009) Cold signalling and cold acclimation in plants. *Advances in Botanical Research* **49**, 35-150.
- Sakai, A. (1978) Low temperature exotherms of winter buds of hardy conifers. *Plant Cell Physiology* **19**, 1439-1446.
- Sakai A. and Larcher W. (1987) Frost Survival of Plants. Responses and Adaptation to Freezing Stress. Springer-Verlag, New York.
- Sakai, A., FuchJgaml, L. and Wetser, C.J. (1986) Cold hardiness in the genu *Rhododendron*. *Journal of the American Society for Horticultural Science* **111**, 273-280.
- Sanders, D., Brownlee, C. and Harper, J.F. (1999) Communicating with calcium. *The Plant Cell* **11**, 691-706.
- Sanghera, G.S., Wani, S.H., Hussain, W. and Singh, N.B. (2011) Engineering cold stress tolerance in crop plants. *Current Genomics* **12**, 30-43.
- Santos, H.P. and Buckeridge, M.S. (2004) The role of the storage carbon of cotyledons in the establishment of seedlings of *Hymenaea courbaril* under different light conditions. *Annals of Botany* **94**, 819-830.
- Schatzberg, P. (1967) On the molecular diameter of water from solubility and diffusion measurements. *The Journal of Physical Chemistry* **71**, 4569-4570.
- Schiefelbein, J.W., Shipley, A. and Rowse, P. (1992) Calcium influx at the tip of growing root-hair cells of *Arabidopsis thaliana*. *Planta* **187**, 455-459.
- Schröder, R., Clark, C.J., Sharrock, K., Hallett, I.C., MacRae, E.A. (2004) Pectins from the

- albedo of immature leon fruitlets have high water binding capacity. *Journal of Plant Physiology* **161**(4), 371-379.
- Schroeder, J.I., Kwak, J.M. and Allen, G.J. (2001) Guard cell abscisic acid signalling and engineering drought hardiness in plants. *Nature* **410**, 327-330.
- Seifert, G.J. and K. Roberts. (2007) The biology of arabinogalactan proteins. *Annual Review of Plant Biology* **58**, 137-161.
- Seki, M., Narusaka, M., Ishida, J., Nanjo, T., Fujita, M., Oono, Y., Kamiya, A., Nakajima, M., Enju, A. and Sakurai, T. (2002) Monitoring the expression profiles of 7000 Arabidopsis genes under drought, cold and high-salinity stresses using a full-length cDNA microarray. *The Plant Journal* **31**, 279-292.
- Sellberg, J.A., Huang, C., McQueen, T.A., Loh, N.D., Laksmono, H., Schlesinger, D., Sierra, R.G., Nordlund, D., Hampton, C.Y., Starodub, D., DePonte, D.P., Beye, M., Chen, C., Martin, A.V., Barty, A., Wikfeldt, K.T., Weiss, T.M., Caronna, C., Feldkamp, J., Skinner, L.B., Seibert, M.M., Messerschmidt, M., Williams, G.J., Boutet, S., Pettersson, L.G.M., Bogan M.J. and Nilsson, A. (2014) Ultrafast X-ray probing of water structure below the homogeneous ice nucleation temperature. *Nature* **510**, 381-384.
- Shevell, D.E., Leu, W.M., Gillmor, C.S., Xia, G., Feldmann, K.A. and Chua, N.H. (1994) *EMB30* is essential for normal cell division, cell expansion and cell adhesion in *Arabidopsis* and encodes a protein that has similarity to Sec7. *Cell* **77**, 1051-1062.
- Shi, H., Ye, T., Zhong, B., Liu, X., Jin, R. and Chan, Z. (2014) AtHAP5A modulates freezing stress resistance in *Arabidopsis* through binding to CCAAT motif of AtXTH21. *New Phytologist* **203**, 554-567.
- Single, W.V. and Marcellos, H. (1981) Ice formation and freezing injury in actively growing cereals. In: *Analysis and Improvement of Plant Cold Hardiness*, Olien, C. R. and Smith, M. N. (eds.), pp. 17-33. CRC Press, Boca Raton, Florida.
- Sivaguru, M., Fujiwara, T., Samaj, J., Baluska, F., Yang, Z.M., Osawa, H., Maeda, T., Mori, T., Volkmann, D. and Matsumoto, H. (2000) Aluminum-induced 1- to 3- $\beta$ -D-glucan inhibits cell-to-cell trafficking of molecules through plasmodesmata: a new mechanism of aluminum toxicity in plants. *Plant Physiology* **124**, 991-1005.
- Solecka, D., Żebrowski, J. and Kacperska, A. (2008) Are pectins involved in cold acclimation and de-acclimation of winter oil-seed rape plants? *Annals of Botany* **101**, 521-530.

- Štajner, D., Milić, N., Čanadanović–Brunet, J., Kapor, A., Štajner, M. and Popović, B.M. (2006) Exploring *Allium* species as a source of potential medicinal agents. *Pytotherapy Research* **20**, 581-584.
- Stefanowska, M., Kuraś, M., Kubacka-Zębalska, M. and Kacperska, A. (1999) Low temperature affects of growth and structure of cell walls in winter oilseed rape (*Brassica napus* L. var. *oleifera* L.) plants. *Annals of Botany* **84**, 313-319.
- Steponkus, P.L. (1990) Cold acclimation and freezing injury from a perspective of the plasma membrane. In: *Environmental Injury to Plants*, Katterman, E. (ed.), pp. 1-16. Academic Press, San Diego, CA.
- Steponkus, P.L., Dowgert, M.F. and Gordon-Kamm, W.J. (1983) Destabilization of the plasma membrane of isolated plant protoplasts during a freeze-thaw cycle: the influence of cold acclimation. *Cryobiology* **20**, 448-65.
- Steponkus, P.L., Dowgert, M.F., Evans, R.Y. and Gordon-Kamm, W. (1982) Cryobiology of isolated protoplasts, pp. 459-474 in Li, P. H. and Sakai, A. (eds.), *Plant Cold Hardiness and Freezing Stress*, Academic Press, New York.
- Steponkus, P.L., Uemura, M., Joseph, R.A., Gilmour, S.J. and Thomashow, M.F. (1998). Mode of action of the COR15a gene on the freezing tolerance of *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences of the United States of America* **95**, 14570-14575.
- Storey, K. and Tanino, K.K. (2012) Temperature adaptation in a changing climate: nature at risk (CABI Climate Change Series). CABI, Oxford, UK .
- Stushnoff, C., Fowler, D.B. and Brule-Babel, A. (1984) Breeding and selection for resistance to low temperature. In: *Crop Breeding: a Contemporary Basis*, Vose, P. B. and Blixt, S. G. (eds.), pp. 115-136. Pergamon Press, Oxford, UK.
- Suutarinen, J., Honkapää, K., Heiniö, R.L., Autio, K. and Morkkila, M. (2000) The effect of different prefreezing treatments on the structure of strawberries before and after jam making. *Lebensmittel-Wissenschaft Und-Technologie-Food Science and Technology* **33**, 188-201.
- Swindell, W.R. (2006) The association among gene expression responses to nine abiotic stress treatments in *Arabidopsis thaliana*. *Genetics* **174**, 1811-1824.

- Taiz, L. (1984) Plant cell expansion: regulation of cell wall mechanical properties. *Annual Review of Plant Physiology* **35**, 585-657.
- Tanino, K.K., Chen, T.H.H., Fuchigami, L.H. and Weiser, C.J. (1990) Metabolic alterations associated with abscisic acid-induced frost hardiness in bromegrass suspension culture cells. *Plant Cell Physiology* **31**, 505-511.
- Tanino, K.K., Chen, T.H.H., Fuchigami, L.H. and Weiser, C.J. (1991) Abscisic acid-induced cellular alterations during the induction of freezing tolerance in bromegrass cells. *Journal of Plant Physiology* **137**, 619-624.
- Tanino, K.K., Kalcsits, L., Silim, S., Kendall, E. and Gray, G. (2010) Temperature-driven plasticity in growth cessation and dormancy development in deciduous woody plants: a working hypothesis suggesting how molecular and cellular function is affected by temperature during dormancy induction. *Plant Molecular Biology* **73**, 49-65.
- Tanino, K.K., Kobayashi, S., Hyett, C., Hamilton, K., Liu, J., Li, B., Borondics, F., Pedersen, T., Tse, J., Ellis, T., Kawamura, Y. and Uemura, M. (2013) *Allium fistulosum* as a novel system to investigate mechanisms of freezing resistance. *Physiologia Plantarum* **147**, 101-111.
- Tao, D., Li, P.H. and Carter, J.V. (1983) Role of cell wall in freezing tolerance of cultured potato cells and their protoplasts. *Physiologia Plantarum* **58**, 527-532.
- The *Arabidopsis* Genome Initiative (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* **408**, 796-815.
- Thomashow, M.F. (1999) Plant cold acclimation: freezing tolerance genes and regulatory mechanisms. *Annual Review of Plant Physiology and Plant Molecular Biology* **50**, 571-599.
- Thomashow, M.F. (2001) So what's new in the field of plant cold acclimation? Lots! *Plant Physiology* **125**, 89-93.
- Thompson, S. (1995) *The Kitchen Garden*. Bantam Books, New York.
- Toivio-Kinnucan, M.A. and Stushnoff, C. (1981) Lipid participation in intracellular freezing avoidance mechanism of lettuce seed. *Cryobiology* **18**, 72-78.
- Trewavas, A.J. and Malho, R. (1998)  $\text{Ca}^{2+}$  signaling in plant cells: The big network! *Current Opinion in Plant Biology* **1**, 428-433.

- Trewavas, A.J. and Gilroy, S. (1991) Signal transduction in plant cells. *Trends in Genetics* **7**, 356-361.
- Tuteja, N. and Mahajan, S. (2007) Calcium signaling network in plants. *Plants Signaling and Behavior* **2**, 79-85.
- Uddin, M.N., Hanstein, S., Leubner, R. and Schubert, S. (2013) Leaf cell-wall components as influenced in the first phase of salt stress in three maize (*Zea mays* L.) hybrids differing in salt resistance. *Journal of Agronomy and Crop Science* **199**, 405-415.
- Uemura, M. and Steponkus, P.L. (1997) Artificial manipulation of the intracellular sucrose content alters the incidence of freeze-induced membrane lesions of isolated protoplasts of *Arabidopsis thaliana*. *Cryobiology* **35**, 336
- Uemura, M., Tominaga, Y., Nakagawara, C., Shigematsu, S., Minami, A. and Kawamura, Y. (2006). Responses of the plasma membrane to low temperatures. *Physiologia Plantarum* **126**, 81-89.
- Valluru, R. and Van den Ende, W. (2008) Plant fructans in stress environments: emerging concepts and future prospects. *Journal of Experimental Botany* **59**, 2905–2916.
- van der Schoot, C. and Rinne, P.L.H. (2011) Dormancy cycling at the shoot apical meristem: Transitioning between self-organization and self-arrest. *Plant Science* **180**, 120-131.
- Verhertbruggen, Y., Marcus, S.E., Haeger, A., Ordaz-Ortiz, J.J. and Knox, J.P. (2009) An extended set of monoclonal antibodies to pectic homogalacturonan. *Carbohydrate Research* **344**, 1858-1862.
- Verslues, P.E., Agarwal, M., Katiyar-Agarwal, S., Zhu, J.H. and Zhu, J.K. (2006) Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *The Plant Journal* **45**, 523-539.
- Vincken, J.P., Schols, H.A., Oomen, R. J.F.J., McCann, M.C., Ulvskov, P., Voragen, A.G.J. and Visser, R.G.F. (2003) If homogalacturonan were a side chain of rhamnogalacturonan I. Implications for cell wall architecture. *Plant Physiology* **132**, 1781-1789.
- Vogel, J. (2008) Unique aspects of the grass cell wall. *Current Opinion in Plant Biology* **11**, 301-307.
- Vogel, J., Raab, T.K., Schiff, C. and Somerville S.C. (2002) PMR6, a pectate lyase-like gene required for powdery mildew susceptibility in Arabidopsis. *The Plant Cell* **14**, 2095-2106.

- Volpi, C., Janni, M., Lionetti, V., Bellincampi, D., Favaron, F. and D'Ovidio, R. (2011) The ectopic expression of a pectin methyl esterase inhibitor increases pectin methyl esterification and limits fungal diseases in wheat. *Molecular Plant-Microbe Interactions Journal* **24**, 1012-1019.
- Vonnegut, B. (1947) The nucleation of ice formation by silver iodide. *Journal of Applied Physics* **18**, 593-594.
- Voragen, A.G.J., Coenen, G.J., Verhoef, R.P. and Schols H.A. (2009) Pectin, a versatile polysaccharide present in plant cell walls. *Structural Chemistry* **20**, 263-275.
- Vorwerk, S., Somerville, S. and Somerville, C. (2004) The role of plant cell wall polysaccharide composition in disease resistance. *Trends in Plant Science* **9**, 203-209.
- Wakeley, P.R., Rogers, H.J., Rozycka, M., Greenland, A.J. and Hussey, P.J. (1998) A maize pectin methylesterase-like gene, *ZmC5*, specifically expressed in pollen. *Plant Molecular Biology* **37**, 187-192.
- Wang, Y., Yang Z.M., Zhang Q. F. and Li J.L. (2009) Enhanced chilling tolerance in *Zoysia matrella* by pre-treatment with salicylic acid, calcium chloride, hydrogen peroxide or 6-benzylaminopurine. *Biologia Plantarum* **53**, 179-182
- Wang, N. and Fisher, D.B. (1994) The use of fluorescent tracers to characterize the post-phloem transport pathway in maternal tissues of developing wheat grains. *Plant Physiology* **104**, 17-28.
- Wang, X., Li, W., Li, M., Welti, R. (2006) Profiling lipid changes in plant response to low temperatures. *Physiologia Plantarum* **126**, 90-96.
- Weiser, C.J. (1970) Cold resistance and injury in woody plants: Knowledge of hardy plant adaptations to freezing stress may help us to reduce winter damage. *Science* **169**, 1269-1278.
- Weiser, R.L., Wallner, S. J. and Waddell, J.W. (1990) Cell wall and extension mRNA changes during cold acclimation of pea seedlings. *Plant Physiology* **93**, 1021-1026.
- Welling, A. and Palva, E.T. (2006) Molecular control of cold acclimation in trees. *Physiologia Plantarum* **127**, 167-181.
- Wenkert, W. (1980) Measurement of tissue osmotic pressure. *Plant Physiology* **65**, 614- 617.
- White, P.J. (1998) Calcium channels in the plasma membrane of root cells. *Annals of Botany* **81**, 173-183.



- White, P.J. and Broadley, M.R. (2003) Calcium in plants. *Annals of Botany* **92**, 487-511.
- Willats, W.G.T., McCartney, L., Mackie, W. and Knox, J.P. (2001) Pectin: cell biology and prospects for functional analysis. *Plant Molecular Biology* **47**, 9-27.
- Wilson, R.H., Smith, A.C., Kačuráková, M., Saunders, P.K., Wellner, N., and Waldron, K.W. (2000) The mechanical properties and molecular dynamics of plant cell wall polysaccharides studied by fourier-transform infrared spectroscopy. *Plant Physiology* **124**, 397-405.
- Wisniewski, M. and Davis, G. (1989) Evidence for the involvement of a specific cell wall layer in the regulation of deep supercooling of xylem parenchyma. *Plant Physiology* **139**, 105-116.
- Wisniewski, M. and Davis, G. (1995) Immunogold localization of pectins and glycoproteins in tissues of peach with reference to deep supercooling. *Trees* **9**, 253-260.
- Wisniewski, M. and Fuller, M. (1999) Ice nucleation and deep supercooling in plants: New insights using infrared thermography. In: *Cold Adapted Organisms: Ecology, Physiology, Enzymology and Molecular Biology*, Margesin, R. and Schinner, F. (eds.). Springer-Verlag, Berlin.
- Wisniewski, M., Ashworth, E. and Schaffer, K. (1987) The use of lanthanum to characterize cell wall permeability in relation to deep supercooling and extracellular freezing in woody plants: II. Intrageneric comparisons between *Betula lenta* and *Betula papyrifera*. *Protoplasma* **141**, 160-168.
- Wisniewski, M., Davis, G. and Arora, R. (1991a) Effect of macerage, oxalic acid, and EGTA on deep supercooling and pit membrane structure of xylem parenchyma of peach. *Plant Physiology* **96**, 1354-1359.
- Wisniewski, M., Davis, G. and Schaffer, K. (1991b) Mediation of deep supercooling of peach and dogwood by enzymatic modifications in cell-wall structure. *Planta* **184**, 254-260.
- Wisniewski, M., Fuller, M., Glenn, D.M., Gusta, L., Duman, J. and Griffith, M. (2002) Extrinsic ice nucleation in plants: what are the factors involved and can they be manipulated. In: *Plant Cold Hardiness: Gene Regulation and Genetic Engineering*, Li, P. H. and Palva, E. T. (eds.), pp.211-221. Springer, US.
- Wisniewski, M., Lindow, S.E. and Ashworth, E.N. (1997) Observation of ice nucleation and propagation in plants using infrared video thermography. *Plant Physiology* **113**, 327-334.

- Wisniewski, M., Nassuth, A., Teulières, C., Marque, C., Rowland, J., Cao, P-B. and Brown, A. (2014) Genomics of cold hardiness in woody plants. *Critical Reviews in Plant Sciences* **33**, 92–124.
- Wolf, S., Mouille, G. and Pelloux, J. (2009) Homogalacturonan methyl-esterification and plant development. *Molecular Plant* **2**, 851-860.
- Wolkers, W.F., McCready, S., Brandt, W.F., Lindsey, G.G. and Hoekstra F.A. (2001). Isolation and characterization of a D-7 LEA protein from pollen that stabilizes glasses *in vitro*. *Biochimica et Biophysica Acta* **1544**, 196-206.
- Wolpert, J.A. and Howell, G.S. (1984) Effect of cane length and dormant season pruning date on cold hardiness and water content of concord bud and cane tissues. *American Journal of Enology and Viticulture* **35**, 237-241.
- Workmaster, B.A.A., Palta, J.P. and Wisniewski, M. (1999) Ice nucleation and propagation in cranberry uprights and fruit using infrared video thermography. *Journal of the American Society for Horticultural Science* **124**, 619-625.
- Wydra, K. and Berl, H. (2006) Structural changes of homogalacturonan, rhamnogalacturonan I and arabinogalactan protein in xylem cell walls of tomato genotypes in reaction to *Ralstonia solanacearum*. *Physiological and Molecular Plant Pathology* **68**, 41-50.
- Wyn Jones, R.G., and Lunt, O.R. (1967). The function of calcium in plants. *Botanical Review* **33**, 407-426.
- Xin, Z. and Browse, J. (1998) *eskimo1* mutants of *Arabidopsis* are constitutively freezing-tolerant. *Plant Biology* **95**, 7799-7804.
- Xiong, L. and Zhu, J.K. (2001) Abiotic stress signal transduction in plants: Molecular and genetic perspectives. *Physiologia Plantarum* **112**, 152-166.
- Xiong, L., Schumaker, K.S. and Zhu, J.K. (2002) Cell signaling during cold, drought, and salt stress. *The Plant Cell* **14**, 165-183.
- Yaish, M.W.F., Doxey, A.C., McConkey, B.J., Moffatt, B.A. and Griffith, M. (2006) Cold-active winter rye glucanases with ice-binding capacity. *Plant Physiology* **141**, 1459-1472.
- Yamasaki, A., Tanaka, K. and Nakashima, N. (2003) Effect of photoperiod on the induction of devernialization by high day temperature in field-grown Japanese bunching onion (*Allium fistulosum* L.). *Journal of the Japanese Society for Horticultural Science* **72**, 18-23.

- Yamauchi, Y., Ogawa, M., Kuwahara, A., Hanada, A., Kamiya, Y. and Yamaguchi, S. (2004) Activation of gibberellin biosynthesis and response pathways by low temperature during imbibition of *Arabidopsis thaliana* seeds. *The Plant Cell* **16**, 367-378.
- Yamazaki, T., Kawamura, Y. and Uemura, M. (2009) Extracellular freezing-induced mechanical stress and surface area regulation on the plasma membrane in cold-acclimated plant cells. *Plant Signaling and Behavior* **4**, 231-233.
- Yamazaki, T., Kawamura, Y., Minami, A. and Uemura, M. (2008) Calcium-dependent freezing tolerance in *Arabidopsis* involves membrane resealing via Synaptotagmin SYT1. *The Plant Cell* **20**, 3389-3404.
- Yeh, S., Moffatt, B.A., Griffith, M., Xiong, F., Yang, D.S.C., Wiseman, S.B., Sarhan, F., Danyluk, J., Xue, Y.Q., Hew, C.L., Doherty-Kirby, A. and Lajoie, G. (2000) Chitinase genes responsive to cold induce antifreeze proteins in winter cereals. *Plant Physiology* **124**, 1251-1263.
- Yelenosky, G. and Guy, C.L. (1989) Freezing tolerance of citrus, spinach, and petunia leaf tissue: osmotic adjustment and sensitivity to freeze induced cellular dehydration. *Plant Physiology* **89**, 444-451.
- Yokoyama, R. and Nishitani, K. (2004) Genomic basis for cell-wall diversity in plants. A comparative approach to gene families in rice and *Arabidopsis*. *Plant and Cell Physiology* **45(9)**, 1111-1121.
- Yoo, S.H., Fishman, M.L., Savary, B.J. and Hotchkiss, A.T.J. (2003) Monovalent salt-induced gelation of enzymatically deesterified pectin. *Journal of Agricultural and Food Chemistry* **51**, 7410-7417.
- Yooyongwech, S., Horigane, A.K., Yoshida, M., Yamaguchi, M., Sekozawa, Y., Sugaya, S. and Gemma, H. (2008) Changes in aquaporin expression and magnetic resonance imaging of water status in peach tree flower buds during dormancy. *Physiologia Plantarum* **134**, 522-533.
- Yu, S.S., Yin, L.K. and Mu, S.Y. (2010) Discovery of an antifreeze protein in the leaves of *Ammopiptanthus nanus*. *Canadian Journal of Plant Science* **90**, 35-40.
- Yun, H.S., Kwon, C., Kim, T.W., Joo, S.H., Cho, M.H., Kang, B.G., Chang, S.C. and Kim, S.K. (2007) Regulation of *VrXTH1* expression in mungbean. *Journal of Plant Biology* **50**: 65-69.

- Zabackis, E., Jing, H., Muller, B., Darvill, A.G., Albersheim, P. (1995) Characterization of the cell-wall polysaccharides of *Arabidopsis thaliana* leaves. *Plant Physiology* **107**, 1129-1138.
- Zabotin, A.I., Barisheva, T.S., Zabolina, O.A., Larskaya, I.A., Lozovaya, V.V., Beldman, G. and Voragen, A.G.J. (1998) Alterations in cell walls of winter wheat roots during low temperature acclimation. *Journal of Plant Physiology* **152**, 473-479.
- Zámečník, J., Bieblová, J. and Grospietsch, M. (1994) Safety zone as a barrier to root-shoot ice propagation. *Plant and Soil* **167**, 149-155.
- Zandleven, J.S., Beldman, G., Bosveld, M., Schols, H.A. and Voragen, A.G.J. (2006) Enzymatic degradation studies of xylogalacturonans from apple and potato, using xylogalacturonan hydrolase. *Carbohydrate Polymers* **65**, 495-503.
- Zeuthen, T. (2001) How water molecules pass through aquaporins. *Trends in Biochemical Sciences* **26**, 77-79.
- Zhang, Z., Nadezhina, E. and Wilkinson, K.J. (2011) Quantifying diffusion in a biofilm of *streptococcus mutans*. *Antimicrobial Agents and Chemotherapy* **55(3)**, 1075-1081.
- Zhou, B. and Guo, Z. (2009) Calcium is involved in the abscisic acid-induced ascorbate peroxidase, superoxide dismutase and chilling resistance in *Stylosanthes guianensis*. *Biologia Plantarum* **53**, 63-68.
- Zhu, H., Derksen, R.C., Krause, C.R., Fox, R.D., Brazee, R.D. and Ozkan, H.E. (2005) Fluorescent intensity of dye solutions under different pH conditions. *Journal of ASTM International* **2**, 1-7.
- Zhu, J.-K. (2002) Salt and drought stress signal transduction in plants. *Annual Review of Plant Biology* **53**, 247-273.
- Zhu, Y-N., Shi, D-Q., Ruan, M-B., Zhang, L-L. , Meng, Z-H., Liu, J. and Yang, W-C. (2013) Transcriptome analysis reveals crosstalk of responsive genes to multiple abiotic stresses in cotton (*Gossypium hirsutum* L.). *Pubic Library of Science One* **8**, e80218.

APPENDIX A: Growth and morphology of *Allium fistulosum* L.



*Allium fistulosum* L. growing in soilless mix in the College of Agriculture and Bioresources greenhouse at University of Saskatchewan



The leaf number sequence of *Allium fistulosum* L.




e: epidermal ligule layer

n: non-epidermal layer


w: whole sheath

The single epidermal layer of *Allium fistulosum* L.

## APPENDIX B: The cryostage microscope equipment




The whole view of cryostage microscope



Upper: Nikon camera attachment  
Lower: Linkam PE94 controller



Heat sink unit



Linkam freezing stage, LTS 120

## APPENDIX C: Protoplasmic streaming and ice penetration in the cells



media1.wmv Live cells with protoplasmic streaming



media2.wmv Dead cells without protoplasmic streaming



media3.wmv Ice penetration into non-acclimated cells

APPENDIX D: Analysis of freezing behavior in treated epidermal layer cells of ligules

The cryostage behavior of NA, ACC and NA+Ca

Treatment <sup>(1)</sup>	AgI extracellular nucleating agent <sup>(2)</sup>	Freezing Temperature (°C)		Viability <sup>(3)</sup>
		extracellular	intracellular	
NA	-	-7.4	no	alive
NA	-	-7.6	no	alive
NA	-	-5.3	no	alive
ACC	-	-9.4	no	alive
ACC	-	-11	no	alive
NA	-	-7.3	-7.4	dead
NA	-	-7.5	-7.5	dead
NA	-	-7.4	-7.4	dead
NA	-	-10.5	-10.6	dead
NA	-	-8.2	-8.2	dead
NA	-	-8.5	-8.5	dead
NA+Ca	-	-10.5	-10.5	dead
NA+Ca	-	-7.3	-7.4	dead
NA+Ca	-	-8.5	-8.6	dead
NA+Ca	-	-11.9	-11.9	dead
ACC	-	-10.1	-10.1	dead
ACC	-	-10.4	-17.1	dead
ACC	-	-10.2	-10.2	dead
ACC	-	-12.1	-12.1	dead
ACC	-	-10.8	down to -23 no	dead
ACC	-	-11.7	-11.7	dead
ACC	-	-7.6	-15.9	dead
ACC	-	-13.1	-13.1	dead
ACC	-	-4.10	down to -23 no	dead
NA	+	-3.1	-7.9	dead



<b>NA</b>	+	-7.9	-10.4	dead
<b>NA</b>	+	-2.5	-7.4	dead
<b>NA</b>	+	-2.9	-7.7	dead
<b>NA+Ca</b>	+	-2.5	down to -23 no	dead
<b>NA+Ca</b>	+	-4.2	down to -23 no	dead
<b>NA+Ca</b>	+	-2.5	-16.0	dead
<b>ACC</b>	+	-5.0	one cell -17.1	alive except intracellular frozen cells
<b>ACC</b>	+	-2.5	down to -23 no	dead
<b>ACC</b>	+	-2.8	down to -23 no	dead
<b>ACC</b>	+	-3.2	down to -23 no	dead
<b>ACC</b>	+	-2.5	down to -23 no	dead

---


(1) NA= non-acclimation; NA+Ca= non-acclimation plus 0.05 M CaCl<sub>2</sub>;

ACC= cold acclimation


(2) +/- means added/not added extracellular nucleating agent

(3) Determined by visible protoplasmic streaming

## APPENDIX E: Confocal microscope and fluorescein penetration




ZEISS LSM 510 Confor2 (Department of Biology, University of Saskatchewan)



Nikon confocal microscope (National Hydrology Research Institute, Saskatoon)


 media4.avi fluorescein penetration in non-acclimated cell

 media5.avi fluorescein penetration in acclimated cells

APPENDIX F: Evos digital inverted microscope (Department of Plant Sciences)



## APPENDIX G: Analysis of pectin methyl esterification in ligule epidermal cells



Micrographs of immunolabelling using JIM 5 and JIM 7 antibodies are shown.  
a: non-acclimated sample with JIM 5; b: acclimated sample with JIM 5;  
c: non-acclimated sample with JIM 7; d: acclimated sample with JIM 7.  
Bars: 100  $\mu\text{m}$ .